The effect of drip irrigation and drip fertigation on N₂O and NO emissions, water saving and grain yields in a maize field in the North China Plain

Di Tian^{a,c}, Yuanyuan Zhang^{a,c}, Yujing Mu^{a,b,c,*}, Yizhen Zhou^{a,c}, Chenglong Zhang^{a,c}, Junfeng Liu^{a,c}

^a Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

^b Center for Excellence in Urban Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, 361021, China

^c University of Chinese Academy of Sciences, Beijing 100049, China

HIGHLIGHTS

- Drip irrigation reduced N₂O (13.8%) and NO (16.7%) emissions.
- Drip fertigation reduced N₂O (7.7%) but increased NO (21.7%) emissions.
- Both drip irrigation and drip fertigation significantly increased the maize yields.
- Drip fertigation is important for agricultural development in water shortage area.

GRAPHICAL ABSTRACT

ARTICLE INFO

Article history:

Received 20 June 2016

Received in revised form 20 September 2016

Accepted 20 September 2016

Available online 22 September 2016

Editor: D. Barcelo

Keywords:

Nitrous oxide

Nitric oxide

Drip irrigation

Water saving

Grain yields

ABSTRACT

N₂O and NO emissions, the water usage and grain yields of a maize field in the North China Plain (NCP) under traditional flood irrigation, drip irrigation and drip fertigation were compared. With respect to the flood irrigation treatment, N₂O emissions were reduced by 13.8% in the drip irrigation treatment and 7.7% in the drip fertigation treatment. NO emissions were reduced to 16.7% in the drip irrigation treatment but increased by 21.7% in the drip fertigation treatment. The molar ratios of NO/N₂O within 2 days after each fertilization event were evidently greater from the drip fertigation treatment than from the flood irrigation treatment, indicating that nitrification was more intensive in the drip fertigation treatment than in the treatment of flood irrigation. Compared with the flood irrigation treatment, evident increase of the maize yields in the drip irrigation treatment (28%) and the drip fertigation treatment (3.7%) were found. Although the drip fertigation treatment could evidently increase NO emission, the 40% water reduction in drip fertigation is of great importance for the sustainable development of agriculture in the NCP where water resources are extremely limited. To mitigate NO emissions from agricultural fields in the NCP with drip fertigation, the addition of a nitrification inhibitor combined with N or nitrate fertilizer was recommended.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Atmospheric nitrous oxide (N₂O) plays important roles in global warming (Myhre et al., 2013) and stratospheric ozone (O₃) depletion

* Corresponding author at: Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China.

E-mail address: yjmu@rcees.ac.cn (Y. Mu).

(Ravishankara et al., 2009). Nitrogen oxides (NO_x), which are mainly emitted as nitric oxide (NO), are involved in tropospheric chemistry in photo-oxidant formation (such as O_3 and peroxyacetyl nitrate [PAN]), which is harmful to human health and vegetation (Taylor, 1969; Shepson et al., 1986; Broadmeadow, 1998). Additionally, nitrate formed by photo-oxidation of atmospheric NO_x is a key component of acid rain that causes the acidification and eutrophication of lakes (Bouwman et al., 2002a). Hence, investigating the sources of atmospheric N_2O and NO_x is important for estimating their impacts on the environment.

Agricultural soil has been considered as an important source for atmospheric N_2O and NO (Akiyama et al., 2004). Global annual emissions from fertilized croplands were 0.11–3.3 Tg N_2O -N (Mosier et al., 1998; Bouwman et al., 2002b; Stehfest and Bouwman, 2006) and 8.9–20 Tg NO -N (Davidson and Kingerlee, 1997; Denman et al., 2007). N_2O and NO are by-products of nitrification (del Prado et al., 2006; Yoshihara et al., 2014) and intermediate products of denitrification (Firestone and Davidson, 1989; Ye et al., 1994) which have been found to be greatly increased by nitrogen fertilizer (Sánchez-Martín et al., 2010).

The North China Plain (NCP) has a cultivated land area of 17.95 million hectares, which accounts for 18.6% of the total agricultural area of China (Liu et al., 2001), and it supplies >50% and 33% of the total wheat and maize in China, respectively (Liang et al., 2011). To increase crop yields, excessive nitrogen fertilizer (Zhao et al., 2006; Ju et al., 2009) and a huge volume of ground water (Zhang et al., 2003) have been supplied to the agricultural fields in the NCP. In this area, the typical nitrogen fertilizer rate is currently >500 kg N $\text{ha}^{-1} \text{yr}^{-1}$ for the maize-wheat rotation system, which largely exceeds crop requirements (Cui et al., 2010). In addition to excessive N application, water is another limiting factor for sustainable agricultural development in the NCP (Meng et al., 2012). The NCP is one of the most water-scarce areas in the world, and the groundwater levels persistently declining (Liu et al., 2008) due to overexploitation for agricultural irrigation (Liu et al., 2001; Zhang et al., 2003). Excessive use of N fertilizer combined with overexploitation of the groundwater have resulted in serious environmental problems, such as ground water pollution (Zhang et al., 1996; Liu et al., 2005a; Ju et al., 2006), and increased N_2O and NO emissions (Ju et al., 2009; Liu et al., 2014). Therefore, optimizing fertilization and irrigation management in the NCP is an urgent priority.

Compared with the inefficient flood irrigation method (the prevailing irrigation method in the NCP), drip irrigation has been shown to be an effective technology for saving water and increasing crop yields (Tilman et al., 2002; Singandhupe et al., 2003; Aujla et al., 2007; Ibragimov et al., 2007). Drip irrigation improves irrigation efficiency by providing water precisely to the crop roots and reducing water losses from sources such as surface evaporation, runoff and deep percolation (Li et al., 2003; Singandhupe et al., 2003; Maisiri et al., 2005). Drip irrigation combined with the application of nitrogen fertilizer dissolved in the irrigation water, usually called fertigation, provides nutrients to crops more precisely than broadcast fertilization (the prevailing fertilization method in the NCP), and it has been verified to be an efficient irrigation method for both saving water and reducing N fertilizer (Singandhupe et al., 2003). Therefore, the Chinese government is currently advocating drip irrigation in the water-limited area of the NCP via demonstrations in villages. Because soil moisture is a key factor that affects the production and diffusion of N_2O and NO in the soil, varying the amount of irrigation water and using different irrigation methods can affect emissions of N_2O and NO from soil (Sánchez-Martín et al., 2008). Many studies on N_2O and NO emissions from agricultural fields under furrow irrigation or flood irrigation have been done (e.g., Rochester, 2003; Grace et al., 2010; Qin et al., 2012; Yan et al., 2013; Zhang et al., 2016; Zhou et al., 2016), but only a few investigations have been conducted for N_2O and NO emissions under drip irrigation (Sánchez-Martín et al., 2008; Scheer et al., 2008; Kennedy et al., 2013; Li et al., 2014; Maris et al., 2015). To our best knowledge, only one study investigated N_2O emissions in a wheat field under drip irrigation in the NCP (Wang et al., 2016), and an evident reduction (14.6%) of N_2O

in the drip irrigation treatment compared to flood irrigation was found. However, a short term investigation (from 1 April to 31 May) by Wang et al. (2016) was less representative for evaluating the actual effect of N_2O reduction from a drip-irrigated agricultural field in the NCP.

Increased crop yields and water savings by drip irrigation are most important in the NCP which is the principal crop production area that has the greatest water shortages in China, but N_2O and NO emissions should also be evaluated before drip irrigation is applied in the vast NCP area because of their significant environmental importance.

In this study, N_2O and NO emissions, water usage and the grain yields in a maize field in the NCP were compared under flood irrigation, drip irrigation and drip fertigation to evaluate the comprehensive effect of drip irrigation.

2. Material and methods

2.1. Field experiment

The study was carried out in an agricultural field in Wangdu County ($38^\circ 1' \text{N}$, $115^\circ 5' \text{E}$), Hebei Province, China. Detailed information about the experimental field is in our previous publications (Zhang et al., 2012; Zhou et al., 2016).

Five treatments were compared in an agricultural field with a total area of 390 m^2 divided into thirteen $3 \times 10 \text{ m}^2$ plots: Duplicate plots for control-flood irrigation (CK-F) were subjected to flood irrigation without fertilization; duplicate plots for control-drip irrigation (CK-D) were subjected to drip irrigation without fertilization; three plots for flood irrigation (FI) were subjected to flood irrigation after urea was broadcasted; three plots for drip irrigation (DI) were subjected to drip irrigation after urea was broadcasted, and three plots for drip fertigation (DF) were subjected to drip fertigation by dissolving urea in the irrigation water. The amounts of urea and irrigation water for each treatment are listed in Table 1. The amounts of urea fertilization and irrigation water in the drip irrigation treatment were the same as for flood irrigation. Compared with flood irrigation, the amounts of urea and water were respectively reduced by 30% and 40% in the drip fertigation treatment. Calcium superphosphate and KCl were broadcast on the flood irrigation, drip irrigation and drip fertigation treated plots. Two fertilization events were conducted for the summer maize based on the cultivating practices of local farmers (Table 1). All of the fertilization treatments were irrigated immediately after fertilization and the control treatments were irrigated on the same days as the fertilization treatments. The summer maize was sowed on July 1, 2013. The maize rows were spaced 50 cm apart, with a distance of 30 cm between the plants in each row. In the drip irrigation and drip fertigation treatments, the distance of every two drip pipes was 50 cm, and the interval between two emitters in each drip pipe was 30 cm. Precipitation and temperature data are shown in Fig. 1 and were obtained from the website <http://www.wunderground.com/>.

CK-F: Control-Flood irrigation; CK-D: Control-Drip irrigation; DI: Drip irrigation; FI: Flood irrigation; DF: Drip fertigation.

Table 1
Application amounts of N in urea and irrigation water in the five treatments.

Date	N-Urea (kg N ha^{-1})	Irrigation (mm)							
		FI	DI	DF	CK-F	CK-D	DI	FI	DF
2013/7/1	Basal fertilization ^a	188	188	131	40	40	40	40	24
2013/8/12	Topdressing ^b	69	69	48	20	20	20	20	12
Total		257	257	179	60	60	60	60	36

^a The basal fertilizers were urea (N = 46%), calcium superphosphate (P_2O_5 = 18%) and KCl (K_2O = 60%) with N: P_2O_5 : K_2O = 25%: 10%: 10%.

^b The fertilizers in the topdressing were urea (N = 46%) and KCl (K_2O = 60%) with N: K_2O = 30%: 5%.

Fig. 1. Daily air temperature (°C) and precipitation (mm) during the period of the field experiment.

2.2. Measurement of the N_2O and NO fluxes

Static chambers were used to measure the N_2O and NO fluxes, and each chamber (I.D. of 32 cm, 80 cm in height) consisted of a cylinder supported by a stainless steel bracket and covered with a Teflon film. Stainless steel pedestals were inserted 10 cm into the soil during the entire maize growing season. N_2O and NO samples were taken by mini-pumps (NMP 830 KNDC, Germany) at 10-min intervals (0, 10, 20 and 30 min) and 3-min intervals (0, 3, 6, 9 min), respectively, after the chambers were deployed on the pedestals. The gas sampling was collected from 8:00 a.m. to 11:00 a.m. on each sampling day. The field measurements were conducted from 29 June to 11 October 2013. Daily sampling of both N_2O and NO lasted for approximately 10 days after each fertilization event, and occurred twice a week for N_2O and occasionally for NO on the other days.

The N_2O concentrations in the air samples were analyzed with a gas chromatograph (GC, Model SP3410, Beijing Analytical Instrument Factory, China) with an electron capture detector (ECD). The temperatures for the GC oven, injector port and ECD were 72 °C, 72 °C and 390 °C, respectively. The NO concentrations were analyzed by a chemiluminescent NH_3 analyzer (Thermo Electron model 17i, USA). The analysis methods for N_2O and NO have been reported in detail in our previous studies (Zhang et al., 2011, 2013).

2.3. Soil sampling and analyzing

Topsoil (0–5 cm) samples were taken with a ring sampler (100 cm³) for soil moisture detection. A gravimetric measurement was done by drying the soil at 105 °C for 24 h. The soil water-filled pore space (WFPS) was calculated according to the following formulas:

$$WFPS(\%) = \frac{\text{Volumetric water content} \times 100}{\text{Soil porosity}}, \quad (1)$$

$$\text{Soil porosity} = \frac{1 - \text{Soil bulk density}}{2.65} \times 100\%. \quad (2)$$

where 2.65 (g cm⁻³) is the theoretical particle density of the soil.

2.4. Data analysis

The N_2O and NO fluxes (F , $\mu\text{g N m}^{-2} \text{h}^{-1}$) were calculated according to the following formula:

$$F = H \times \frac{\Delta C}{\Delta t} \times \frac{P}{RT} \times M. \quad (3)$$

where H (m) is the height of the chamber, $\Delta C/\Delta t$ is the slope of the linear regression of the N_2O and NO concentrations (C , ppbV) versus time (t , h), P (atm) is the atmospheric pressure measured in the field, R

(0.082 atm L/(K · mol)) is the gas constant, T (K) is the ambient air temperature and M (g/mol) is the molecular weight of nitrogen.

The emission factor (EF, %) was calculated based on the following formula:

$$EF = \frac{E_N - E_0}{N} \times 100. \quad (4)$$

where E_N and E_0 are the amounts of the cumulative N_2O -N or NO-N emissions (kg N ha^{-1}) from the fertilized plots and the control plots, respectively. N is the total N (kg N ha^{-1}) input.

The data were analyzed using SPSS 22.0 software (SPSS Inc., Chicago, USA) and plotted with Origin Pro 8.5 (Origin Lab Corporation, USA). The cumulative N_2O -N emissions and NO-N emissions were estimated by successive linear interpolation based on the sampling date and the assumption that the emissions followed a linear trend over the periods when no samples were taken. The significance of comparisons of the grain yields, the emission factors and the cumulative emissions between different treatments were determined by LSD method ($P < 0.05$) in a one-way ANOVA analysis.

3. Results and discussion

3.1. N_2O emissions

The variations of the N_2O emissions from the five treatments during the entire maize growing season are shown in Fig. 2. Compared with the two control treatments, notable N_2O emissions from the three fertilization treatments primarily occurred within approximately 10 days after each fertilization event and were in good agreement with the results of our previous studies in the same field (Zhang et al., 2011, 2012, 2014; Zhou et al., 2016). An evident difference in N_2O emissions was present among the three fertilization treatments. After the basal fertilization, the N_2O emission peaks were $1948.8 \mu\text{g N m}^{-2} \text{h}^{-1}$ for the flood irrigation treatment, $2322.3 \mu\text{g N m}^{-2} \text{h}^{-1}$ for the drip fertigation treatment and $1677.6 \mu\text{g N m}^{-2} \text{h}^{-1}$ for the drip irrigation treatment, and appeared on the first day, the second day and the third day, respectively. Although the amount of N fertilizer applied in the drip fertigation treatment was only 70% of that in the flood irrigation and drip irrigation treatments, the N_2O emission peak from the drip fertigation treatment was the highest of the three treatments after the basal fertilization, which was ascribed to the different fertilization procedures and differences in soil moisture. Urea was applied in the drip fertigation treatment by dripping a water solution, but it was broadcasted into the flood irrigation and drip irrigation treatments. Because granular urea could not be completely dissolved in the flood and drip irrigation treatments, only part of the urea was involved in N_2O

Fig. 2. Variations of the N_2O emissions from the control-flood irrigation, control-drip irrigation, flood irrigation, drip irrigation and drip fertigation treatments during the maize growing period. The solid arrows indicate fertilization and irrigation. Error bars represent the standard deviation.

production in those treatments after irrigation, which was confirmed by the additional N_2O emission peaks from the two treatments on 8–9 July (Fig. 2) just after a strong rainfall (~75 mm, Fig. 1). In addition, N_2O production and diffusion in the soil strongly depends on soil moisture (Firestone and Davidson, 1989; Russow et al., 2008), and the relatively lower soil moisture in the drip fertigation treatment (Fig. 3) might have favored N_2O production and diffusion.

In contrast to the basal fertilization, the N_2O emission peaks from the three fertilization treatments were more narrowed and appeared on the same day (the third day) after the topdressing. The relatively narrow N_2O emission peaks were probably due to the rapid N utilization by the maize during the topdressing period via the competition with the N utilization from microbial processes (Inselsbacher et al., 2010; Kennedy et al., 2013). The basal fertilizer was applied to the field just after the maize was seeded, hence almost all of the N fertilizer was available for N_2O production from soil nitrification and denitrification, resulting in a relatively long period of high N_2O emissions. Compared with the cumulative N_2O emissions from the two fertilization events (basal fertilization and topdressing), the N_2O emissions from the agricultural field could possibly be reduced if the basal fertilization was postponed until the maize leaves emerged. The soil moisture might be less disturbed by irrigation due to the high frequency of rainfall during the topdressing period (Fig. 1 and Fig. 3), so the N_2O emission peaks from the three fertilization treatments appeared on the same day after the topdressing and irrigation. During the topdressing period, the N_2O emission peak from the flood irrigation treatment ($1861.7 \mu\text{g N m}^{-2} \text{h}^{-1}$) was approximately 2-fold higher than those from the drip irrigation treatment ($996.0 \mu\text{g N m}^{-2} \text{h}^{-1}$) and the drip fertigation treatment ($1023.0 \mu\text{g N m}^{-2} \text{h}^{-1}$). Compared with the flood irrigation treatment, the notable reduction of N_2O emissions in both drip irrigation and drip fertigation treatments after topdressing indicated that drip irrigation favored of the utilization nitrogenous fertilizer by the crop during the growing season (Wu et al., 2014a).

3.2. NO emissions

The variations of NO emissions from the five treatments during the entire maize growing season are shown in Fig. 4. Similar to the N_2O emissions, the NO emissions from the three fertilization treatments immediately increased and lasted for about ten days after each fertilization event. However, the NO emission peaks after the basal fertilization event were postponed for 1–2 days compared with the corresponding N_2O emission peaks, which was in agreement with our previous studies (Zhang et al., 2016; Zhou et al., 2016). NO emission from soil was the net effect of its production and diffusion rates (Russow et al., 2008; Oswald et al., 2013), and thus the earlier emission peaks of N_2O than those of NO were rationally ascribed to a strong restriction of NO diffusion in the soil under extremely high moisture conditions that were present immediately after irrigation. Although the irrigation water and the amount of

Fig. 3. The soil water-filled pore space (WFPS, %) at a 0–5 cm depth from the flood irrigation, drip irrigation and drip fertigation treatments during the maize season. Dashed arrows indicate irrigation only.

Fig. 4. Variations of the NO emissions from the control-flood irrigation, control-drip irrigation, flood irrigation, drip irrigation and drip fertigation treatments during the maize growing period. Solid arrows show fertilizer application and irrigation. Error bars represent the standard deviation.

fertilizer in the drip fertigation treatment were the lowest of the three treatments, the highest emission peak (Jul 4, $1046.4 \mu\text{g N m}^{-2} \text{h}^{-1}$) was observed from drip fertigation. Compared with the basal fertilization, NO emissions from the three fertilization treatments after topdressing were much lower, which was in good agreement with the results of previous studies (Hosono et al., 2006; Zhang et al., 2011; Zhou et al., 2016). The relatively low NO emissions after topdressing were ascribed to the relatively high soil moisture, a lower amount of N fertilizer application and the rapid utilization of nitrogenous fertilizer by the maize crop.

The NO emissions from the agricultural fields were mainly ascribed to nitrification, whereas the N_2O emissions were from both nitrification and denitrification (Skiba et al., 1993), hence the molar ratio of $\text{NO}/\text{N}_2\text{O}$ has been used as a useful indicator for distinguishing the relative contributions from soil nitrification and denitrification (Anderson and Levine, 1986; Williams et al., 1998; Meijide et al., 2007). A laboratory simulation indicated that the molar ratio of $\text{NO}/\text{N}_2\text{O}$ was usually greater than one for nitrifiers and much less than one for denitrifiers (Anderson and Levine, 1986). However, the field measurements revealed that the nitrification predominantly occurred when $\text{NO}/\text{N}_2\text{O} > 0.11$ (Meijide et al., 2007). As shown in Fig. 5, the molar ratios of $\text{NO}/\text{N}_2\text{O}$ from both flood irrigation and drip fertigation were >0.11 within 7 days after the basal fertilization, and 2 days after the topdressing. The molar ratios of $\text{NO}/\text{N}_2\text{O}$ were much greater for the drip fertigation treatment than for the flood irrigation within 2 days after the basal fertilization and topdressing, respectively. The sum NO emissions during the 2 days accounted for 33–36% and 69–100% of the cumulative emissions (within 9 days after each fertilization event) from the flood irrigation and the drip fertigation, respectively. Therefore, the relatively high ratio of $\text{NO}/\text{N}_2\text{O}$ and the high proportion of NO emissions from the drip fertigation treatment occurred within the 2 days after each fertilization event, implying that nitrification was the predominant source of the NO and N_2O emissions from the treatment. Because the optimum soil

Fig. 5. Molar ratios of $\text{NO}/\text{N}_2\text{O}$ from the flood irrigation and drip fertigation treatments during the maize growing period. Solid arrows show fertilizer application and irrigation.

moisture (WFPS) is usually below 60% for nitrification and above 60% for denitrification (Davidson et al., 1991; Bateman and Baggs, 2005; del Prado et al., 2006), the relatively low soil moisture in the drip fertigation treatment (Fig. 3) compared with the flood irrigation treatment might favor nitrification, resulting in relatively high NO emission peaks. On the other hand, the soil moisture also affects NO diffusion from the soil to the atmosphere – the lower the soil moisture is, the faster NO diffusion rate (Liu et al., 2005b; Russow et al., 2008). Additionally, the granular urea could not be completely dissolved in the flood and drip irrigations (as previously mentioned), so only part of the urea was initially involved in NO production after irrigation, whereas the nutrient N in the water solution of urea in the drip fertigation treatment could be effectively used by the microbes for nitrification.

3.3. The grain yields, the emission factors and cumulative emissions

The maize yields, emission factors and cumulative emissions of N₂O and NO for the different treatments are listed in Table 2. The maize yields from drip irrigation and drip fertigation treatments were significantly increased by 28% and 3.7%, respectively, with respect to the flood irrigation treatment.

The emission factors of N₂O and NO were lower for the drip irrigation treatment than for the flood irrigation treatment, whereas they were significantly greater for the drip fertigation treatment than for the flood irrigation treatment. The N₂O emission factors for the three treatments in this study were in the range of 0.15%–3.8% as reported by other studies for maize fields in the NCP (Meng et al., 2005; Cui et al., 2012; Liu et al., 2014), whereas the NO emission factors were lower than 0.52% as reported by Cui et al. (2012) and 0.76% reported by Zhang et al. (2011).

Compared with the flood irrigation treatment, the cumulative N₂O and NO emissions per unit area from the drip irrigation treatment were decreased by 13.8% and 16.7%, respectively. The cumulative N₂O and NO emissions per unit of yield were significantly reduced in the drip irrigation treatment to 32.7% and 35.3%, respectively. In contrast to the reduction of cumulative N₂O emissions from the drip fertigation treatment, the cumulative NO emissions from the drip fertigation treatment were significantly greater than those from the flood irrigation treatment.

Wang et al. (2016) reported that the reduction percentage of N₂O emissions was 14.6% in a wheat field in the NCP under drip irrigation with respect to flood irrigation, which was in good agreement with the result obtained in this study. The reduction percentages of the N₂O emissions from the melon and cotton fields under drip irrigation with

respect to furrow irrigation could be as high as 28%–70% (Sánchez-Martín et al., 2008, 2010; Wu et al., 2014a). The large differences in the N₂O reduction percentages for drip irrigation in various studies might be due to the different origins of the N₂O emissions from the fields. N₂O emissions from an agricultural field under flood irrigation in the NCP has been verified to be predominantly due to nitrification (Ju et al., 2011; Zhang et al., 2011; Cui et al., 2012), whereas denitrification was found to be the predominant process for N₂O emissions from agricultural fields under furrow irrigation (Sánchez-Martín et al., 2008, 2010; Aguilera et al., 2013).

The evident increase of NO emissions for the drip fertigation treatment with respect to flood and drip irrigation treatments further confirmed that a relatively dry soil favored nitrification and NO diffusion (Davidson et al., 2000). Sánchez-Martín et al. (2010) also found that NO emissions from an agricultural field increased with the application of organic fertilizer under drip irrigation compared with furrow irrigation. However, Sánchez-Martín et al. (2008) also reported that the NO emissions were reduced to 33% from the same agricultural field with application of ammonium sulfate under drip irrigation compared with furrow irrigation. To explain these inconsistent results, further field comparison studies are still required.

3.4. The importance of the application of drip irrigation and drip fertigation in the NCP

Maize fields occupy 10.18 million ha (Wu et al., 2014b), and account for 51% of the cultivated land area in the NCP. If the traditional flood irrigation was completely replaced by the drip irrigation in the NCP as done in this study, the reduction in the amount of N₂O during the maize season would be 9.76×10^6 kg, which accounts for 4.23% of the total emissions (2.31×10^8 kg N₂O yr⁻¹, Zhao et al., 2012; Zhou et al., 2014) in this region. Based on the results of this study, the reduction of the NO_x emissions (1.02×10^6 kg N) from the maize fields of the NCP with 100% replacement of traditional flood irrigation by drip irrigation only accounts for 0.10% of the total NO_x emissions (1.08×10^9 kg N yr⁻¹) from the entire region in the NCP, 0.30% of the NO_x emissions (0.34×10^9 kg N yr⁻¹) from the traffic sector, and 0.67% of the NO_x emission (0.15×10^9 kg N yr⁻¹) from industrial processing (Zhao et al., 2012).

Assuming that drip fertigation with a 40% water savings was completely applied in the NCP, the reduction in the amount of N₂O during the maize season would be 5.44×10^6 kg which accounts for 2.35% of the total emissions in this region. The application of drip fertigation during the maize season would also save 2.44×10^9 m³ of water, which

Table 2

The grain yields (water content, 15.5%), the emission factors and the cumulative emissions of N₂O and NO (mean ± SE) during summer maize growing season.

	CK-F	CK-D	FI	DI	DF ^a
Yield					
Grain yield (kg ha ⁻¹)	4531 ± 160a	6622 ± 161b	8841 ± 121c	11,320 ± 72e	9171 ± 78d
N ₂ O ^b					
EF (%)	–	–	1.57 ± 0.11ab	1.33 ± 0.13a	2.05 ± 0.62b
Emission (kg N ₂ O-N ha ⁻¹)	0.39 ± 0.09a	0.41 ± 0.26a	4.43 ± 0.28b	3.82 ± 0.34b	4.09 ± 1.12b
Emission/yield (mg N ₂ O-N kg ⁻¹)	86.55 ± 19.32a	62.16 ± 38.72a	500.88 ± 31.67c	337.19 ± 29.75b	446.29 ± 121.99bc
NO ^c					
EF (%)	–	–	0.24 ± 0.03a	0.19 ± 0.03a	0.39 ± 0.03b
Emission (kg NO-N ha ⁻¹)	–0.0013 ± 0.002a	0.025 ± 0.001a	0.60 ± 0.08b	0.50 ± 0.08b	0.73 ± 0.05c
Emission/yield (mg NO-N kg ⁻¹)	–0.29 ± 0.45a	3.72 ± 0.18a	68.18 ± 8.98c	44.13 ± 7.34b	79.88 ± 5.92c

CK-F: Control-Flood irrigation; CK-D: Control-Drip irrigation; DI: Drip irrigation; FI: Flood irrigation; DF: Drip fertigation.

Different letters within columns denote significant differences by applying the Least Significant Difference (LSD) test at $P < 0.05$.

^a The emission factor from the DF treatment was calculated based on the emission from the CK-D treatment.

^b N₂O emissions is the cumulative emissions during the entire maize growing season.

^c NO emissions is the cumulative emissions during approximately 10 days after fertilization.

accounts for 3.47% of the total water use ($7.05 \times 10^{10} \text{ m}^3 \text{ yr}^{-1}$, NBSC, 2015) and 5.78% of the agricultural water use ($4.23 \times 10^{10} \text{ m}^3 \text{ yr}^{-1}$, NBSC, 2015). If drip fertigation was adopted during the whole year for the typical winter wheat–summer maize rotation system, the ground-water savings potential would be greater because >70% of the irrigation water is applied to wheat in the NCP (Li et al., 2005). Additionally, the application of the drip fertigation in the NCP maize fields would reduce N fertilizer use $7.9 \times 10^8 \text{ kg N yr}^{-1}$, which accounts for 14.03% of the total N fertilizer use in the NCP ($5.63 \times 10^9 \text{ kg N yr}^{-1}$, NBSC, 2015).

Because photochemical O_3 formation is usually limited by NO_x in rural areas (NRC, 1992; Stohl et al., 1996), the increase of NO emissions from the drip fertigation would exacerbate O_3 formation if traditional flood irrigation was replaced by the drip fertigation in the NCP. Application of a nitrification inhibitor and nitrate fertilizer have been found to be effective control measures for both N_2O and NO emissions from agricultural fields in the NCP (Ding et al., 2010; Ju et al., 2011; Hu et al., 2013; Zhang et al., 2016; Zhou et al., 2016), therefore we recommend a nitrification inhibitor combined with N fertilizer and nitrate fertilizer to mitigate NO emissions from the agricultural fields in the NCP under drip fertigation.

4. Conclusions

Drip irrigation could decrease N_2O and NO emissions from the NCP maize fields compared to the current practice of flood irrigation, but an increase of NO emissions was found with the drip fertigation. Compared with flood irrigation, both drip irrigation and drip fertigation could significantly increase maize yields. Drip fertigation using a nitrification inhibitor combined with N fertilizer or nitrate fertilizer is recommended to mitigate NO emissions from the agricultural fields in the NCP and would be beneficial for protecting the local atmosphere and for the sustainable development of agriculture in regions with severe water shortages.

Acknowledgements

This work was supported by the National Key Technology R&D Program of China (No. 2013BAD11B03-3), the Strategic Priority Research Program of the Chinese Academy of Sciences (No. XDB05010100), and the National Natural Science Foundation of China (Nos. 41575121, 91544211 and 21477142).

References

- Aguilera, E., Lassaletta, L., Sanz-Cobena, A., Garnier, J., Vallejo, A., 2013. The potential of organic fertilizers and water management to reduce N_2O emissions in Mediterranean climate cropping systems. A review. *Agric. Ecosyst. Environ.* 164, 32–52.
- Akiyama, H., McTaggart, I.P., Ball, B.C., Scott, A., 2004. N_2O , NO, and NH_3 emissions from soil after the application of organic fertilizers, urea and water. *Water Air Soil Poll.* 156, 113–129.
- Anderson, I.C., Levine, J.S., 1986. Relative rates of nitric oxide and nitrous oxide production by nitrifiers, denitrifiers, and nitrate respirers. *Appl. Environ. Microbiol.* 51 (5), 938–945.
- Aujla, M.S., Thind, H.S., Buttar, G.S., 2007. Fruit yield and water use efficiency of eggplant (*Solanum melongena* L.) as influenced by different quantities of nitrogen and water applied through drip and furrow irrigation. *Sci. Hortic.* 112, 142–148.
- Bateman, E.J., Baggs, E.M., 2005. Contributions of nitrification and denitrification to N_2O emissions from soils at different water-filled pore space. *Biol. Fert. Soils* 41, 379–388.
- Bouwman, A.F., Boumans, L.J.M., Batjes, N.H., 2002a. Emissions of N_2O and NO from fertilized fields: Summary of available measurement data. *Glob. Biogeochem. Cycl.* 16 (4), 6–16–13.
- Bouwman, A.F., Boumans, L.J.M., Batjes, N.H., 2002b. Modeling global annual N_2O and NO emissions from fertilized fields. *Glob. Biogeochem. Cycl.* 16 (4), 28–21–28–29.
- Broadmeadow, M., 1998. Ozone and forest trees. *New Phytol.* 139, 123–125.
- Cui, F., Yan, G., Zhou, Z., Zheng, X., Deng, J., 2012. Annual emissions of nitrous oxide and nitric oxide from a wheat–maize cropping system on a silt loam calcareous soil in the North China Plain. *Soil Biol. Biochem.* 48, 10–19.
- Cui, Z., Chen, X., Zhang, F., 2010. Current nitrogen management status and measures to improve the intensive wheat–maize system in China. *Ambio* 39, 376–384.
- Davidson, E.A., Kingery, W., 1997. A global inventory of nitric oxide emissions from soils. *Nutr. Cycl. Agroecosyst.* 48, 37–50.
- Davidson, E.A., Rogers, J.E., Whitman, W.B., 1991. Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems. In: Rogers, J.E., Whitman, W.B. (Eds.), *Microbial Production and Consumption of Greenhouse Gases: Methane, Nitrogen Oxides, and Halomethanes*, pp. 219–235.
- Davidson, E.A., Keller, M., Erickson, H.E., Verchot, L.V., Veldkamp, E., 2000. Testing a conceptual model of soil emissions of nitrous and nitric oxides. *Bioscience* 50, 667–680.
- Denman, K.L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P.M., Dickinson, R.E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P.L., Wofsy, S.C., Zhang, X., 2007. Couplings between changes in the climate system and biogeochemistry. In: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (Eds.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Ding, W.X., Yu, H.Y., Cai, Z.C., 2010. Impact of urease and nitrification inhibitors on nitrous oxide emissions from fluvo-aquic soil in the North China plain. *Biol. Fert. Soils* 47, 91–99.
- Firestone, M.K., Davidson, E.A., 1989. Microbiological basis of NO and N_2O production and consumption in soil. In: Andreae, M.O., Schimel, D.S. (Eds.), *Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere*. Wiley, Chichester, pp. 7–21.
- Grace, P., Rowlings, D., Rochester, I., Kiese, R., Butterbach-Bahl, K., Gilkes, R.J., Prakongkep, N., 2010. Nitrous oxide emissions from irrigated cotton soils of northern Australia. 19th World Congress of Soil Science, Soil Solutions for a Changing World, 1–6 August 2010 Brisbane, Australia Publ. DVD, pp. 179–182.
- Hosono, T., Hosoi, N., Akiyama, H., Tsuruta, H., 2006. Measurements of N_2O and NO emissions during tomato cultivation using a flow-through chamber system in a glasshouse. *Nutr. Cycl. Agroecosyst.* 75, 115–134.
- Hu, X.-K., Su, F., Ju, X.-T., Gao, B., Oenema, O., Christie, P., Huang, B.-X., Jiang, R.-F., Zhang, F.-S., 2013. Greenhouse gas emissions from a wheat–maize double cropping system with different nitrogen fertilization regimes. *Environ. Pollut.* 176, 198–207.
- Ibragimov, N., Evett, S.R., Esanbekov, Y., Kamilov, B.S., Mirzaev, L., Lamers, J.P.A., 2007. Water use efficiency of irrigated cotton in Uzbekistan under drip and furrow irrigation. *Agric. Water Manag.* 90, 112–120.
- Inselsbacher, E., Hinko-Najera Umana, N., Stange, F.C., Gorfer, M., Schüller, E., Ripka, K., Zechmeister-Boltenstern, S., Hood-Novotny, R., Strauss, J., Wanek, W., 2010. Short-term competition between crop plants and soil microbes for inorganic N fertilizer. *Soil Biol. Biochem.* 42, 360–372.
- Ju, X.T., Kou, C.L., Zhang, F.S., Christie, P., 2006. Nitrogen balance and groundwater nitrate contamination: comparison among three intensive cropping systems on the North China Plain. *Environ. Pollut.* 143, 117–125.
- Ju, X.T., Xing, G.X., Chen, X.P., Zhang, S.L., Zhang, L.J., Liu, X.J., Cui, Z.L., Yin, B., Christie, P., Zhu, Z.L., Zhang, F.S., 2009. Reducing environmental risk by improving N management in intensive Chinese agricultural systems. *Proc. Natl. Acad. Sci. U. S. A.* 106, 3041–3046.
- Ju, X.T., Lu, X., Gao, Z., Chen, X.P., Su, F., Kogge, M., Romheld, V., Christie, P., Zhang, F.S., 2011. Processes and factors controlling N_2O production in an intensively managed low carbon calcareous soil under sub-humid monsoon conditions. *Environ. Pollut.* 159, 1007–1016.
- Kennedy, T.L., Suddick, E.C., Six, J., 2013. Reduced nitrous oxide emissions and increased yields in California tomato cropping systems under drip irrigation and fertigation. *Agric. Ecosyst. Environ.* 170, 16–27.
- Li, J., Zhang, J., Ren, L., 2003. Water and nitrogen distribution as affected by fertigation of ammonium nitrate from a point source. *Irrig. Sci.* 22, 19–30.
- Li, J., Inanaga, S., Li, Z., Eneji, A.E., 2005. Optimizing irrigation scheduling for winter wheat in the North China Plain. *Agric. Water Manag.* 76, 8–23.
- Li, Z., Zhang, R., Wang, X., Chen, F., Lai, D., Tian, C., 2014. Effects of plastic film mulching with drip irrigation on N_2O and CH_4 emissions from cotton fields in arid land. *J. Agric. Sci.* 152, 534–542.
- Liang, Q., Chen, H., Gong, Y., Fan, M., Yang, H., Lal, R., Kuzyakov, Y., 2011. Effects of 15 years of manure and inorganic fertilizers on soil organic carbon fractions in a wheat–maize system in the North China Plain. *Nutr. Cycl. Agroecosyst.* 92, 21–33.
- Liu, C., Yu, J., Eloise, K., 2001. Groundwater exploitation and its impact on the environment in the North China Plain. *Water Int.* 26, 265–272.
- Liu, C., Yao, Z., Wang, K., Zheng, X., 2014. Three-year measurements of nitrous oxide emissions from cotton and wheat–maize rotational cropping systems. *Atmos. Environ.* 96, 201–208.
- Liu, G.D., Wu, W.L., Zhang, J., 2005a. Regional differentiation of non-point source pollution of agriculture-derived nitrate nitrogen in groundwater in northern China. *Agric. Ecosyst. Environ.* 107, 211–220.
- Liu, J., Zheng, C., Zheng, L., Lei, Y., 2008. Ground water sustainability: methodology and application to the North China Plain. *Ground Water* 46, 897–909.
- Liu, X.J., Mosier, A.R., Halvorson, A.D., Zhang, F.S., 2005b. Tillage and nitrogen application effects on nitrous and nitric oxide emissions from irrigated corn fields. *Plant Soil* 276, 235–249.
- Maisiri, N., Senzanje, A., Rockstrom, J., Twomlow, S.J., 2005. On farm evaluation of the effect of low cost drip irrigation on water and crop productivity compared to conventional surface irrigation system. *Phys. Chem. Earth A, B, C* 30, 783–791.
- Maris, S.C., Teira-Esmatges, M.R., Arbones, A., Rufat, J., 2015. Effect of irrigation, nitrogen application, and a nitrification inhibitor on nitrous oxide, carbon dioxide and methane emissions from an olive (*Olea europaea* L.) orchard. *Sci. Total Environ.* 538, 966–978.
- Meijide, A., Díez, J.A., Sánchez-Martín, L., López-Fernández, S., Vallejo, A., 2007. Nitrogen oxide emissions from an irrigated maize crop amended with treated pig slurries and composts in a Mediterranean climate. *Agric. Ecosyst. Environ.* 121, 383–394.
- Meng, L., Ding, W., Cai, Z., 2005. Long-term application of organic manure and nitrogen fertilizer on N_2O emissions, soil quality and crop production in a sandy loam soil. *Soil Biol. Biochem.* 37, 2037–2045.
- Meng, Q., Sun, Q., Chen, X., Cui, Z., Yue, S., Zhang, F., Röhmheld, V., 2012. Alternative cropping systems for sustainable water and nitrogen use in the North China Plain. *Agric. Ecosyst. Environ.* 146, 93–102.
- Mosier, A., Kroeze, C., Nevison, C., Oenema, O., Seitzinger, S., Clempu, O.V., 1998. Closing the global atmospheric N_2O budget: nitrous oxide emissions through the agricultural nitrogen cycle. *Nutr. Cycl. Agroecosyst.* 33, 3538–3543.

- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, B.M., Nakajima, T., Robock, A., Stephens, G., Takemura, T., Zhang, H., 2013. Anthropogenic and natural radiative forcing. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- NBSC (National Bureau of Statistic of China), 2015. *China Statistical Year Book*. China Statistics Press, Beijing.
- NRC (National Research Council), 1992. *Rethinking the ozone problem in urban and regional air pollution*. Commission on Geosciences, Environment and Resources. National Academy of Sciences Press, Washington, DC.
- Oswald, R., Behrendt, T., Ermel, M., Wu, D., Su, H., Cheng, Y., Breuninger, C., Moravek, A., Mougín, E., Delon, C., 2013. HONO emissions from soil bacteria as a major source of atmospheric reactive nitrogen. *Science* 341, 1233–1235.
- del Prado, A., Merino, P., Estavillo, J.M., Pinto, M., González-Murua, C., 2006. N₂O and NO emissions from different N sources and under a range of soil water contents. *Nutr. Cycl. Agroecosyst.* 74, 229–243.
- Qin, S., Wang, Y., Hu, C., Oenema, O., Li, X., Zhang, Y., Dong, W., 2012. Yield-scaled N₂O emissions in a winter wheat–summer corn double-cropping system. *Atmos. Environ.* 55, 240–244.
- Ravishankara, A., Daniel, J.S., Portmann, R.W., 2009. Nitrous oxide (N₂O): the dominant ozone-depleting substance emitted in the 21st century. *Science* 326, 123–125.
- Rochester, I.J., 2003. Estimating nitrous oxide emissions from flood-irrigated alkaline grey clays. *Soil Res.* 41, 197–206.
- Russow, R., Spott, O., Stange, C.F., 2008. Evaluation of nitrate and ammonium as sources of NO and N₂O emissions from black earth soils (Haplic Chernozem) based on ¹⁵N field experiments. *Soil Biol. Biochem.* 40, 380–391.
- Sánchez-Martín, L., Arce, A., Benito, A., García-Torres, L., Vallejo, A., 2008. Influence of drip and furrow irrigation systems on nitrogen oxide emissions from a horticultural crop. *Soil Biol. Biochem.* 40, 1698–1706.
- Sánchez-Martín, L., Meijide, A., García-Torres, L., Vallejo, A., 2010. Combination of drip irrigation and organic fertilizer for mitigating emissions of nitrogen oxides in semiarid climate. *Agric. Ecosyst. Environ.* 137, 99–107.
- Scheer, C., Wassmann, R., Klenzler, K., Lbragimov, N., Eschanov, R., 2008. Nitrous oxide emissions from fertilized irrigated cotton (*Gossypium hirsutum* L.) in the Aral Sea Basin, Uzbekistan: influence of nitrogen applications and irrigation practices. *Soil Biol. Biochem.* 40, 290–301.
- Shepson, P.B., Kleindienst, T.E., Edney, E.O., Nero, C.M., Cupitt, L.T., Claxton, L.D., 1986. Acetaldehyde: the mutagenic activity of its photooxidation products. *Environ. Sci. Technol.* 20, 1008–1013.
- Singandhupe, R.B., Rao, G.G.S.N., Patil, N.G., Brahmanand, P.S., 2003. Fertigation studies and irrigation scheduling in drip irrigation system in tomato crop (*Lycopersicon esculentum* L.). *Eur. J. Agronomy* 19, 327–340.
- Skiba, U., Smith, K.A., Fowler, D., 1993. Nitrification and denitrification as sources of nitric oxide and nitrous oxide in a sandy loam soil. *Soil Biol. Biochem.* 25, 1527–1536.
- Stehfest, E., Bouwman, L., 2006. N₂O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions. *Nutr. Cycl. Agroecosyst.* 74, 207–228.
- Stohl, A., Williams, E., Wotawa, G., Kromp-Kolb, H., 1996. A European inventory of soil nitric oxide emissions and the effect of these emissions on the photochemical formation of ozone. *Atmos. Environ.* 30, 3741–3755.
- Taylor, O.C., 1969. Importance of peroxyacetyl nitrate (PAN) as a phytotoxic air pollutant. *J. Air Pollut. Control. Assoc.* 19, 347–351.
- Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural sustainability and intensive production practices. *Nature* 418, 671–677.
- Wang, G.S., Liang, Y.P., Zhang, Q., Jha, S.K., Gao, Y., Shen, X.J., Sun, J.S., Duan, A.W., 2016. Mitigated CH₄ and N₂O emissions and improved irrigation water use efficiency in winter wheat field with surface drip irrigation in the North China Plain. *Agric. Water Manag.* 163, 403–407.
- Williams, P.H., Jarvis, S.C., Dixon, E., 1998. Emission of nitric oxide and nitrous oxide from soil under field and laboratory conditions. *Soil Biol. Biochem.* 30, 1885–1893.
- Wu, J., Guo, W., Feng, J.F., Li, L.H., Yang, H.S., Wang, X.H., Bian, X.M., 2014a. Greenhouse gas emissions from cotton field under different irrigation methods and fertilization regimes in arid northwestern China. *Sci. World J.* 407832.
- Wu, L., Chen, X., Cui, Z., Zhang, W., Zhang, F., 2014b. Establishing a regional nitrogen management approach to mitigate greenhouse gas emission intensity from intensive smallholder maize production. *PLoS One* 9, 22.
- Yan, G., Zheng, X., Cui, F., Yao, Z., Zhou, Z., Deng, J., Xu, Y., 2013. Two-year simultaneous records of N₂O and NO fluxes from a farmed cropland in the northern China plain with a reduced nitrogen addition rate by one-third. *Agric. Ecosyst. Environ.* 178, 39–50.
- Ye, R.W., Averill, B.A., Tiedje, J.M., 1994. Denitrification: production and consumption of nitric oxide. *Appl. Environ. Microbiol.* 60, 1053–1058.
- Yoshihara, T., Tokura, A., Hashida, S.-N., Kitazaki, K., Asobe, M., Enbutsu, K., Takenouchi, H., Goto, F., Shoji, K., 2014. A precise/short-interval measurement of nitrous oxide emission from a rockwool tomato culture. *Environ. Control. Biol.* 52, 137–147.
- Zhang, W., Tian, Z., Zhang, N., Li, X., 1996. Nitrate pollution of groundwater in northern China. *Agric. Ecosyst. Environ.* 59, 223–231.
- Zhang, X., Pei, D., Hu, C., 2003. Conserving groundwater for irrigation in the North China Plain. *Irrigation Sci.* 21, 159–166.
- Zhang, Y., Liu, J., Mu, Y., Pei, S., Lun, X., Chai, F., 2011. Emissions of nitrous oxide, nitrogen oxides and ammonia from a maize field in the North China Plain. *Atmos. Environ.* 45, 2956–2961.
- Zhang, Y., Liu, J., Mu, Y., Xu, Z., Pei, S., Lun, X., Zhang, Y., 2012. Nitrous oxide emissions from a maize field during two consecutive growing seasons in the North China Plain. *J. Environ. Sci.* 24, 160–168.
- Zhang, Y., Mu, Y., Fang, S., Liu, J., 2013. An improved GC-ECD method for measuring atmospheric N₂O. *J. Environ. Sci.* 25, 547–553.
- Zhang, Y., Mu, Y., Zhou, Y., Liu, J., Zhang, C., 2014. Nitrous oxide emissions from maize-wheat field during 4 successive years in the North China Plain. *Biogeosciences* 11, 1717–1726.
- Zhang, Y., Mu, Y., Zhou, Y., Tian, D., Liu, J., Zhang, C., 2016. NO and N₂O emissions from agricultural fields in the North China Plain: origination and mitigation. *Sci. Total Environ.* 551–552, 197–204.
- Zhao, B., Wang, P., Ma, J.Z., Zhu, S., Pozzer, A., Li, W., 2012. A high-resolution emission inventory of primary pollutants for the Huabei region, China. *Atmos. Chem. Phys.* 12, 481–501.
- Zhao, R., Chen, X., Zhang, F., Zhang, H., Schroder, J., Römhild, V., 2006. Fertilization and nitrogen balance in a wheat–maize rotation system in north China. *Agron. J.* 98, 938–945.
- Zhou, F., Shang, Z., Ciais, P., Tao, S., Piao, S., Raymond, P., He, C., Li, B., Wang, R., Wang, X., Peng, S., Zeng, Z., Chen, H., Ying, N., Hou, X., Xu, P., 2014. A new high-resolution N₂O emission inventory for China in 2008. *Environ. Sci. Technol.* 48, 8538–8547.
- Zhou, Y., Zhang, Y., Tian, D., Mu, Y., 2016. Impact of dicyandiamide on emissions of nitrous oxide, nitric oxide and ammonia from agricultural field in the North China Plain. *J. Environ. Sci.* 40, 20–27.