

Arsenic Accumulation and Speciation in Maize as Affected
by Inoculation with Arbuscular Mycorrhizal Fungus
Glomus mosseae

YANG YU, SHUZHEN ZHANG,* HONGLIN HUANG, LEI LUO, AND BEI WEN

State Key Laboratory of Environmental Chemistry and Ecotoxicology, Research Center for
Eco-Environmental Sciences, Chinese Academy of Sciences, P.O. Box 2871, Beijing 100085, China

Effects of inoculation with arbuscular mycorrhizal (AM) fungus (*Glomus mosseae*) on arsenic (As) accumulation and speciation in maize were investigated by using As spiked soil at the application levels of 0, 25, 50, and 100 mg kg⁻¹. Inorganic As was the major species in plants, and mycorrhizal inoculation generally decreased concentrations of arsenite [As(III)] in maize roots and concentrations of As(III) and arsenate [As(V)] in the shoots. Dimethylarsenic acid (DMA) concentrations (detected in every plant sample) were higher in maize shoots for mycorrhizal than for nonmycorrhizal treatment, but no significant differences were observed for roots. Monomethylarsenic acid (MMA) was only detected in roots with mycorrhizal colonization. The uptake of As(V) was much lower by excised mycorrhizal than nonmycorrhizal roots, and the differences for the uptake of As(III) were negligible. Arsenate reductase (AR) activity was detected in maize roots, and it was reduced with mycorrhizal inoculation. Activities of peroxidase (POD) and superoxide dismutase (SOD) were detected in both maize shoots and roots, and they were suppressed by mycorrhizal inoculation. AM inoculation inhibited the uptake of As(V) and its reduction to As(III), reducing oxidation stress and thereby alleviating As toxicity to the host plant.

KEYWORDS: Arbuscular mycorrhizal fungus; arsenic speciation; uptake; arsenate reductase activity; antioxidative enzyme activity

INTRODUCTION

Arsenic is a ubiquitous trace metalloid (1) that can be found throughout the world in soils polluted by mining and smelting industries and coal burning, particularly in South East Asia (2). Arsenic in soil can enter the food chain via plant uptake and cause adverse effects on human health (3).

Different As species such as arsenate [As(V)], arsenite [As(III)], monomethylarsenic acid (MAA), and dimethylarsinic acid (DMA) have been found in soils and plants. They have different chemical behavior and are thought to be taken up by roots and transported within plants by different ways. Arsenate acts as a phosphate analogue and is transported across the plasma membrane via phosphate cotransport systems (2). Arsenite has been evidenced to transport into rice roots via glycerol transporting channels (4). Toxic effects of As are highly dependent on its speciation. Inorganic As [As(III) and As(V)] is generally more toxic than the organic species, and As(III) is more toxic than As(V) (1, 2, 5).

Arbuscular mycorrhizal (AM) fungi are ubiquitous in the rhizosphere, forming symbiotic associations with roots of the majority of plant species (6). Their capacity to enhance the tolerance of host plants to As contamination in soil has been addressed (7, 8). AM fungi have also been reported to reduce plant uptake of As such as by the fern *Pteris vittata*

L (9), lentil (10), alfalfa (11), tomato (12), and sunflower (13). Considering the interactions between AM fungi with soil microorganisms in the rhizosphere and their influence on plant uptake of nutrition such as P, we speculated that AM fungi might influence As speciation in soil and plant growth in As contaminated soil. To our knowledge, such information is lacking, and the knowledge is essential for us to understand the effect of AM fungi on the biogeochemistry of As in the soil–plant system.

Reduction and methylation of As can generate reactive oxygen species (ROS) and damage DNA, proteins, and lipids (2). Plants have evolved a variety of mechanisms to alleviate the deleterious effects caused by ROS, generally through the production of various antioxidative enzymes such as superoxide dismutase (SOD) and peroxidase (POD). For example, SOD, a metallo-enzyme, can catalyze the dismutation of O₂⁻ into H₂O₂ and O₂, and subsequently H₂O₂ can be effectively scavenged by POD. A previous study has found that lipid peroxidation and SOD activities are correlated with increasing As(V) concentrations in *Holcus lanatus* L. (14). Srivastava et al. (15) have observed a higher antioxidative enzyme activity in correspondence with As accumulation in As-treated *Pteris vittata*. However, As(V) in most organisms can be enzymatically reduced to As(III) (16). Arsenate reductase (AR) has been detected in terrestrial plants, including ferns, rice, and *Arabidopsis* (17), and has been confirmed to play an important role in plant accumulation and detoxification of As (18).

*Corresponding author. Tel: +86 10 62849683. Fax: +86 10 62923563. E-mail: szzhang@rcees.ac.cn.

The aim of the present study was to investigate the influence of AM inoculation on As accumulation and speciation in maize growing in a soil artificially contaminated with various levels of As(V). The uptake of As(V) and As(III) by excised roots was characterized, and activities of AR and antioxidative enzymes were analyzed and compared between mycorrhizal and nonmycorrhizal plants in order to understand the mechanisms involved in the influence of AM fungus on plant accumulation of different As species and the resistance of mycorrhizal plant to As contamination in soil.

MATERIALS AND METHODS

Inoculum and Soil Preparation. Inoculum of the AM fungus, *Glomus mosseae*, isolated from a noncontaminated soil, was propagated for 10 weeks in pot culture on broomcorn (*Sorghum vulgare* Pers.) plants grown in a soil–sand mixture in a greenhouse. The inoculum, which was air-dried and passed through a 2-mm sieve, consisted of spores, mycelium, sandy soil, and root fragments containing approximately 60 spores g⁻¹ soil (dry weight soil basis).

A sandy soil (Alfisol) was collected from a farm near Beijing. It was dried and passed through a 2-mm sieve. The soil has the following properties (on a dry weight soil basis): pH (1:2.5 soil to water), 8.41; organic matter, 0.39%; 0.5 mol L⁻¹ NaHCO₃-extractable P, 2.75 mg kg⁻¹; and 0.5 mol L⁻¹ NaHCO₃-extractable As, 0.18 mg kg⁻¹. The soil was sterilized by γ -irradiation (10 kGy, 10 MeV γ ray) for the elimination of indigenous AM fungi and received an application of 200 mg kg⁻¹ N (NH₄NO₃), 30 mg kg⁻¹ P (KH₂PO₄), and 150 mg kg⁻¹ K (K₂SO₄) as a basal fertilizer.

Plant Growth and Harvest. Four replicates for each As treatment, 0 mg kg⁻¹, 25 mg kg⁻¹, 50 mg kg⁻¹, and 100 mg kg⁻¹, were applied to mycorrhizal and nonmycorrhizal treatments. Each pot received a mixture of 750 g soil and 50 g inoculum for mycorrhizal treatment or sterilized inoculum plus 15 mL of inoculum washings filtered through a 37 μ m filter paper for nonmycorrhizal treatment. Arsenate was added as Na₂HAsO₄·7H₂O in solution. The pots were allowed to equilibrate for a period of 4 weeks in a greenhouse by undergoing four cycles of saturation with distilled water and air drying. Seeds of maize (*Zea mays* L.) were surface sterilized in a 10% (v/v) solution of hydrogen peroxide and then pregerminated on a moist filter paper for 3 days. Four seedlings were transplanted into each pot, and two uniform seedlings were left after 4 days. The experiment was carried out in a controlled environment glasshouse with a photoperiod of 14 h at a light intensity of 250 μ mol m⁻² s⁻¹ provided by supplementary illumination. The temperature was 25 °C at daytime and 18 °C at night. The moisture content of the soil was maintained at 50% of water holding capacity by regular watering.

Plants were harvested after 8 weeks of growth. Shoots and roots were separately harvested. Root fragments were collected by sieving the soil and adding them to the root samples. Roots were first carefully washed with tap water to remove any adhering soil particles. Then roots and shoots were thoroughly rinsed with distilled water, blotted dry, and weighed. A subsample of fresh roots was taken from each pot, cleared, and stained for the determination of the proportion of root length colonized by the AM fungus and quantified by a grid line intersect method (19). Some fresh roots and shoots were frozen by liquid nitrogen and then stored at -80 °C for enzyme analysis. The remaining plant samples were freeze-dried, ground, and stored at 4 °C, and the dry weights were recorded. The fresh to dry root ratio was used to estimate the total dry mass of roots. Only the rhizosphere soils were sampled from each pot considering the interactions between roots/mycorrhizas and As and the importance of As biogeochemistry in the rhizosphere soil. The soil that required continued, vigorous rubbing and shaking of the root system was classified as rhizosphere soil (20).

Analyses of P, Total As, and As Speciation. Subsamples of 0.1 g dry plant materials were digested with 2.5 mL of concentrated HNO₃ and 2.5 mL of H₂O₂, and 0.1 g of dry soil samples with 2.5 mL of concentrated HNO₃ and 2 mL of concentrated HClO₄. The digestion tubes were heated on a heating block at 120 °C for 1 h and then at 180 °C to evaporate the samples to near dryness. Reagent blank and certified reference material of tea (GBW 08505) were included to verify the accuracy and precision of the digestion procedure, and subsequent analysis and the recovery rates were within 95 \pm 10%. Arsenic was determined by hydride generation atomic fluorescence spectrometry (HG-AFS, AF-610A atomic fluorescence spectrophotometer, Beijing RuiLi Instrumental Company, Beijing, China), and P was determined by the Molybdenum blue method.

Two grams of soil samples or 0.2 g of freeze-dried powdered plant samples were ultrasonically extracted with 10 mL of 1:1 methanol/water for 2 h (21). The samples were centrifuged, and the supernatant was decanted into a polyethylene bottle. The procedure was repeated with the residual pellet, and the two extracts were combined. The extraction solutions were then air-dried with nitrogen to remove methanol and regenerated in a smaller volume (1–5 mL) with double deionized water. The spike recoveries, individual test for As(V), As(III), MMA, and DMA, were within 100 \pm 10%. Extraction efficiency of As, defined as the ratio of As concentration obtained by extraction to that by nitric acid digestion, ranged from 50 to 70% for roots and 75 to 95% for shoots, respectively.

Arsenic speciation in the sample extracts was determined by using HPLC (Agilent 1200 series HPLC) coupled to ICP-MS (Agilent 7500c ICP-MS). An anion exchange column at 150 mm \times 4.6 mm i.d. was used and 2 mM/0.2 mM phosphate buffer/EDTA eluent (pH 6.0) was used to detect As(V), As(III), MMA, and DMA in less than 10 min. One guard column at 10 mm \times 4.6 mm i.d. was used to protect the main column. Each As species in the extracts was verified by the coincidence of retention times with standards. Stock solutions of As(III), DMA, MMA, and As(V) were prepared from NaAsO₂, CH₃AsO(ONa)₂, (CH₃)₂AsO(OH), and Na₂HAsO₄·7H₂O. Detection limits were 0.1, 0.2, 0.1, and 0.1 μ g L⁻¹ for As(III), As(V), DMA, and MMA, respectively. Quality assurance was obtained through the use of blanks, standard curves, standard check solutions, and spiked samples, which were run during sample analysis.

Uptake Kinetics of As(V) and As(III). Plants for the experiment of root uptake of As(V) and As(III) were prepared separately. Methods for inoculation and plant growth in non-arsenic soil were the same as described above. Experimental method for As uptake by roots was adopted from the one by Gonzalez-Chavez et al. (7). Roots of both nonmycorrhizal and mycorrhizal plants were first carefully washed by soaking in water to remove soil particles and excised at the basal node. The excised roots (0.2–0.5 g fresh weight) were incubated in aerated test solution [containing 5.0 mM MES and 0.5 mM Ca(NO₃)₂, adjusted to pH 5 using KOH] for 30 min at 25 °C and then transferred into aerated test solutions containing different concentrations of As(V) or As(III). After the termination of As exposure for 20 min, the roots were rinsed in ice-cold phosphate solution containing 1 mM K₂HPO₄ to remove the absorbed As from the root free space and to stop further root activity. The roots were then oven-dried at 70 °C for 2 days and weighed. Digestion and analysis of the samples were conducted following the methods described above.

Enzyme Analysis. Arsenic reductase (AR) activity was assayed using the coupled enzymatic reaction described by Duan et al. (17) with some modifications. Two grams of fresh plant materials were ground and homogenized to a fine paste in 10 mL of extraction buffer (50 mM MOPS and 50 mM MES, adjusted to pH 6.5 with NaOH), filtered, centrifuged, then purified with Sephadex PD-10 desalting columns. All steps were performed on ice, and the supernatant was stored at 4 °C prior to analysis of AR activity. The assay of AR was performed in 50 mM MOPS and 50 mM MES, pH 6.5, containing 1.5 mM NADPH, 1 unit

yeast glutathione reductase, 1 mM GSH, and 10 mM sodium arsenate. Plant extracts were preincubated for 5 min in the buffer containing glutathione reductase and GSH, and then NADPH and sodium arsenate were added to start the reaction. AR activity was monitored by the decrease of NADPH absorbance at 340 nm, and NADPH oxidation was calculated using a molar extinction coefficient of 6200 for NADPH at 340 nm and expressed in nkat mg⁻¹ protein. One nkat is defined as the amount of enzyme required to oxidize 1 nmol NADPH in 1 s.

A half gram of fresh plant samples were ground and homogenized in 10 mL of solution containing 50 mM potassium phosphate buffer and 1% (w/v) polyvinylpyrrolidone (pH 7.8) on an ice-bath. The homogenate was centrifuged at 8000g for 15 min at 4 °C. The supernatant was stored at 4 °C prior to the analysis of SOD and POD activities, and the soluble protein contents. POD activity was assayed following the method of Polle et al. (22). Changes in the absorbance of brown guaiacol at 460 nm in the presence of H₂O₂ were recorded for calculating POD activity. The enzyme activity was defined as the nmol of H₂O₂ oxidized min⁻¹ mg⁻¹ protein. SOD activity was measured by the spectrophotometric method. The assay was performed in terms of SOD ability to inhibit the reduction of *p*-nitroblue tetrazolium chloride (NBT) to formazan by the superoxide radical. One unit of SOD activity (U) was defined as the amount of enzyme required to inhibit NBT reduction by 50% measured at 560 nm and expressed in U mg⁻¹ protein. Protein contents were determined using Coomassie Brilliant Blue G-250 as dye and bovine serum albumin as the standard. All spectrophotometric analyses were conducted using an ultraviolet–visible spectrophotometer (U-3010, Hitachi Ltd., Tokyo, Japan).

Statistical Analysis. All results were expressed as an average of four replications. Treatment effects were determined by two-way analysis of variance using SPSS 11.0 software.

RESULTS

Colonization Rate, Biomass, and Concentrations of P and As. No mycorrhizal colonization was observed in the roots of noninoculated plants, while mycorrhizal plants had 74, 72, 76, and 69% of root colonization rate at As addition levels of 0, 25, 50, and 100 mg kg⁻¹, respectively, and As concentration in soil did not significantly affect mycorrhizal inoculation ($P > 0.05$).

Table 1 displays the maize biomass, P concentrations, and concentration and total accumulation of As in maize. Both root and shoot biomass decreased markedly when As was applied at 50 and 100 mg kg⁻¹. Mycorrhizal inoculation increased the root and shoot dry weights significantly

($P < 0.001$), especially when As was applied at 100 mg kg⁻¹ where the biomass of mycorrhizal maize was almost twice as much as that of nonmycorrhizal plant. Mycorrhizal inoculation increased P concentration in both roots and shoots at all As application levels ($P < 0.01$). The concentration and total accumulation of As increased consistently in shoots and roots of both nonmycorrhizal and mycorrhizal plants with increasing application rate of As in soil. Mycorrhizal inoculation significantly reduced As concentration in both maize roots and shoots ($P < 0.001$). The total As accumulation in roots and shoots were lower in mycorrhizal than in nonmycorrhizal treatments without As addition or at low As additional level (e.g., 25 mg kg⁻¹), while the opposite pattern was observed when As was applied at high concentration (e.g., 50 and 100 mg kg⁻¹), which could be ascribed to the limited influence of mycorrhizal inoculation on plant biomass at low As addition levels and a significant influence at high As addition levels.

Arsenic Speciation. Inorganic As was the major species found in the rhizosphere soil and plant samples and accounted for more than 97% of the total contents. Concentrations of As(V) and As(III) in maize roots and shoots and the rhizosphere soils after plant harvest are given in **Figure 1**. As(V) and As(III) were detected in all of the samples except that no As(III) was detected in the nonarsenic addition soil. Concentrations of As(V) and As(III) increased consistently in the rhizosphere soils, shoots, and roots with increasing As application rate. In soil, As(V) was the predominant species (91–100%), and mycorrhizal inoculation decreased the concentrations of both As(V) and As(III). Arsenite was the major species present in maize roots (48–85%) and shoots (68–93%). AM inoculated roots accumulated consistently more As(V) and less As(III) than nonmycorrhizal roots. Shoot concentrations of As(V) and As(III) were all obviously lower in the mycorrhizal than in the nonmycorrhizal treatment ($P < 0.001$).

Concentrations of organic As species (MMA and DMA) in the rhizosphere soils and plants are given in **Table 2**. DMA was detected in all of the samples irrespective of As application rates and mycorrhizal treatments with only one exception for the control treatment without As addition to soil, whereas MMA was only found in the rhizosphere soils and roots for nonmycorrhizal treatment. Both MMA and DMA increased with increasing As concentration in the soils. DMA concentrations were observed lower in the soils ($P < 0.05$) and higher in maize shoots ($P < 0.05$) for

Table 1. Dry Biomass, Concentrations of P, and Concentration and Total Accumulation of As in Roots and Shoots of Maize Inoculated with or without AM Fungus (Mean \pm SE, $n = 4$)

As addition (mg kg ⁻¹)	inoculation treatment	dry biomass (g)		P concentration (mg g ⁻¹)		As concentration (mg kg ⁻¹)		total As accumulation (μ g pot ⁻¹)	
		root	shoot	root	shoot	root	shoot	root	shoot
0	NM ^a	4.3 \pm 0.1	7.6 \pm 0.6	0.63 \pm 0.04	0.89 \pm 0.07	1.5 \pm 0.1	0.14 \pm 0.02	6.3 \pm 0.7	1.0 \pm 0.1
	M	4.9 \pm 0.3	9.6 \pm 0.4	1.10 \pm 0.07	1.43 \pm 0.05	1.1 \pm 0.2	0.08 \pm 0.02	5.2 \pm 1.3	0.8 \pm 0.2
25	NM	4.6 \pm 0.3	7.0 \pm 0.1	0.59 \pm 0.11	0.98 \pm 0.06	32.4 \pm 2.4	1.17 \pm 0.13	149.7 \pm 11.6	8.1 \pm 0.7
	M	6.4 \pm 0.2	9.2 \pm 0.3	0.90 \pm 0.04	1.45 \pm 0.03	21.2 \pm 5.1	0.72 \pm 0.13	133.8 \pm 30.8	6.6 \pm 1.3
50	NM	3.6 \pm 0.3	4.4 \pm 0.1	0.61 \pm 0.07	0.80 \pm 0.10	129.2 \pm 17.2	1.56 \pm 0.15	461.7 \pm 37.8	6.8 \pm 0.8
	M	6.1 \pm 0.4	7.8 \pm 0.6	0.95 \pm 0.12	1.34 \pm 0.07	82.1 \pm 10.6	1.04 \pm 0.21	491.6 \pm 26.0	8.2 \pm 2.4
100	NM	2.2 \pm 0.1	2.8 \pm 0.1	0.72 \pm 0.06	0.81 \pm 0.08	175.2 \pm 9.3	2.47 \pm 0.23	381.1 \pm 28.6	6.9 \pm 0.5
	M	4.2 \pm 0.1	6.2 \pm 0.3	1.08 \pm 0.13	1.60 \pm 0.11	121.7 \pm 10.4	1.63 \pm 0.25	505.7 \pm 17.9	10.2 \pm 2.4
significance of inoculation (I)		***	***	***	**	***	***	***	***
arsenic (As)		***	***	NS	NS	***	***	***	***
I \times As		***	***	NS	NS	***	***	***	***

^a NM and M represent nonmycorrhizal and mycorrhizal treatments, respectively. ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$; NS, not significant.

mycorrhizal treatment compared with nonmycorrhizal treatment; however, such differences were not statistically significant for roots ($P > 0.05$).

Effects of Inoculation with AM Fungus on the Uptake Kinetics of As(V) and As(III). The uptake of As(V) and As(III) by maize roots showed a hyperbolic increase with increasing concentration of As in solution (Figure 2), and the uptake kinetics were adequately described by the Michaelis–Menten function (Table 3). The uptake of As(V) was much higher than that of As(III) by both mycorrhizal and nonmycorrhizal roots. A significant difference for the uptake of As(V) existed between mycorrhizal and nonmycorrhizal roots. Uptake of As(V) by mycorrhizal roots had a lower V_{max} value and higher K_m value than that by

nonmycorrhizal roots ($P < 0.05$). But the uptake curves for As(III) by nonmycorrhizal and mycorrhizal roots were almost overlapped, and there were no significant differences between kinetic parameters for mycorrhizal and nonmycorrhizal treatments.

Activities of AR, POD, and SOD. Results of enzyme analysis are shown in Table 4. Arsenate reductase (AR) activities were only detected in roots of maize grown in As spiked soils, and they were progressively increased with increasing As concentration in the soil. Mycorrhizal inoculation significantly decreased AR activities in roots ($P < 0.01$). Both SOD activities in roots and shoots and POD activities in roots consistently increased with increasing As concentration in the soil ($P < 0.001$), while there is no consistent trend in POD activities in shoots. AM inoculation remarkably reduced POD and SOD activities in both roots and shoots ($P < 0.05$).

DISCUSSION

A significant observation of this study was that AM inoculation changed the accumulation of As(V) and As(III) in maize roots and shoots. Mycorrhizal inoculation increased As(V) concentration but decreased As(III) concentration in roots, and both concentrations of As(V) and As(III) in shoots were lower for mycorrhizal compared to those for nonmycorrhizal maize (Figure 1). As(V) and As(III) are taken up by roots and transported through different mechanisms (2, 4), and there is also conversion between As species inside plants (23).




Figure 1. Concentrations of As(V) and As(III) in rhizosphere soils (A), roots (B), and shoots (C) of nonmycorrhizal (open bar) and mycorrhizal (closed bar) maize. Means and standard errors ($n = 4$) on a dry matter basis are presented. I, As, and I \times As represent P values of analysis of variance for the effects of inoculation, arsenic, and their interactions on As speciation, respectively. NS, not significant.




Figure 2. Concentration-dependent kinetics for As(V) and As(III) uptake by maize roots inoculated with or without AM fungus. NM and M represent nonmycorrhizal and mycorrhizal, respectively. Means and standard errors ($n = 4$) on a dry matter basis are presented. Data are the means of four replicates and on a dry matter basis. Bars, standard errors.

Table 2. Concentrations of Organic Arsenic Species in the Rhizosphere Soils, Roots, and Shoots of Maize Inoculated with or without AM Fungus (Mean \pm SE, $n = 4$)

As addition (mg kg ⁻¹)	inoculation treatment	MMA concentrations (μ g kg ⁻¹)			DMA concentrations (μ g kg ⁻¹)		
		soil	root	shoot	soil	root	shoot
0	NM ^a	— ^b	—	—	—	—	—
	M	—	—	—	—	—	—
25	NM	2.0 \pm 0.9	27 \pm 20	—	2.8 \pm 2.0	19 \pm 16	1.5 \pm 0.3
	M	—	—	—	0.8 \pm 0.4	18 \pm 14	1.4 \pm 0.4
50	NM	2.4 \pm 0.4	52 \pm 6	—	2.1 \pm 0.7	61 \pm 19	2.4 \pm 1.1
	M	—	—	—	1.9 \pm 0.7	21 \pm 4	4.9 \pm 1.5
100	NM	12.5 \pm 5.4	106 \pm 68	—	2.9 \pm 1.8	46 \pm 6	13.3 \pm 4.2
	M	—	—	—	1.3 \pm 0.6	40 \pm 13	27.2 \pm 15.2
significance of inoculation (I)					*	NS	*
arsenic (As)					*	***	***
I \times As					NS	NS	NS

^a NM and M represent nonmycorrhizal and mycorrhizal treatments, respectively. ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$; NS, not significant. ^b —, not detected.

We therefore hypothesized that the influence of mycorrhizal inoculation on different accumulation of As(V) and As(III) in maize could come from the different uptake and transport of As(V) and As(III), and/or the different reduction ratios of As(V) to As(III) in mycorrhizal and nonmycorrhizal plants. Therefore, we first conducted an experiment to exam the uptake of As(V) and As(III) by excised mycorrhizal and nonmycorrhizal roots. The results showed that mycorrhizal inoculation remarkably inhibited As(V) uptake but did not significantly influence the uptake of As(III) by roots. Mycorrhizal hyphae can enter into plant cells and directly take up some elements from soil (24). Therefore, beside the effects of inoculation on root uptake of As brought about by morphological and physiological changes in roots, mycorrhizal hyphae might selectively take up and transport different As species, leading to the different accumulations of As(V) and As(III) in maize. Possible translocation difference between As(V) and As(III) inside plants could also result in different accumulations of As(V) and As(III) in roots and shoots. Therefore, the shoot to root ratios of As(V) and As(III) concentrations were calculated (Figure 3) to check whether mycorrhizal inoculation influenced the transport of As(V) and As(III). Mycorrhizal inoculation decreased the ratios of As(V) ($P < 0.01$) but did not affect the ratios of As(III) ($P > 0.05$). On the basis of the combined evidence of limited translocation of As from roots to shoots (data in Table 1) and as well as the limited influence of mycorrhizal inoculation on As translocation from root to shoot (Figure 3), we would conclude that translocation of As(V) and As(III) inside plants was not the key factor determining their different accumulations between mycorrhizal and nonmycorrhizal maize. Furthermore, reduction of As(V) to As(III) might occur inside plants (23), and inoculation could have effects on such reactions, influencing the accumulation of As(V) and As(III) in maize. Up to now, it has been

difficult to trace the conversion between As species inside living plants. AR is a specific catalyzer to reduce As(V) to As(III) in plants (17, 25), and AR activities have been reported to play an important role in As accumulation in plants (18). Therefore, in order to clarify whether mycorrhizal inoculation could inhibit the reduction of As(V) to As(III) in roots, the activities of AR in maize roots were detected. It was found that AR activities were lower in mycorrhizal than in nonmycorrhizal roots. Lower AR activities in mycorrhizal roots indicated that less As(V) was reduced to As(III) compared with nonmycorrhizal roots, which is consistent with the result of lower As(III) concentration in mycorrhizal roots and provides evidence for the influence of mycorrhizal inoculation on the reduction of As(V) to As(III) in maize roots. Furthermore, after inhibition of its reduction by AR, As(V) could be immobilized by mycorrhizal roots as evidenced by the decreased shoot to root ratios of As(V). The role of AR in the reduction reaction and metabolism of As inside plants has recently attracted research attention, but only a small number of studies have been published. It is still unknown whether mycorrhizal inoculation decreases the expression of the AR gene in plants. This study showed the primary evidence that

Table 3. Kinetic Parameters for As(V) and As(III) Influx into Maize Roots Inoculated with or without AM Fungus

As species	inoculation treatment	V_{max} (nmol g ⁻¹ d.wt. h ⁻¹)	K_m (mM)	R^2
As(V)	NM ^a	966 ± 61*	0.0070 ± 0.0022*	0.9313
	M	520 ± 72	0.0313 ± 0.0112	0.8873
As(III)	NM	467 ± 50	0.0534 ± 0.0121	0.9648
	M	604 ± 170	0.1022 ± 0.0483	0.9107

^a NM and M represent nonmycorrhizal and mycorrhizal treatments respectively. *, means significant difference ($P < 0.05$) between nonmycorrhizal and mycorrhizal treatments in each As species.




Figure 3. Shoot to root ratio of As(V) concentration (A) and As(III) concentration (B) in nonmycorrhizal (open bar) and mycorrhizal (closed bar) maize. Means and standard errors ($n = 4$) on a dry matter basis are presented. I, As, and I × As represent P values of analysis of variance for the effects of inoculation, arsenic, and their interactions on As speciation, respectively. NS, not significant.

Table 4. Activities of AR (Arsenate Reductase), POD, and SOD in Roots and Shoots of Maize Inoculated with or without AM Fungus (Mean ± SE, $n = 4$)

As addition (mg kg ⁻¹)	inoculation treatment	AR (nkat mg ⁻¹ protein)		POD (nmol min ⁻¹ mg ⁻¹ protein)		SOD (U mg ⁻¹ protein)	
		root	shoot	root	shoot	root	shoot
0	NM ^a	— ^b	—	559 ± 120	312 ± 40	30.5 ± 4.5	34.5 ± 4.5
	M	—	—	571 ± 31	292 ± 28	36.3 ± 3.5	24.6 ± 4.1
25	NM	9.0 ± 3.2	—	697 ± 96	363 ± 65	41.7 ± 7.7	35.5 ± 7.5
	M	8.3 ± 2.7	—	631 ± 57	240 ± 22	32.4 ± 4.3	29.6 ± 2.7
50	NM	15.5 ± 2.3	—	936 ± 74	315 ± 25	63.1 ± 6.3	44.8 ± 3.5
	M	11.6 ± 1.1	—	725 ± 67	270 ± 16	46.8 ± 3.6	35.6 ± 2.1
100	NM	20.9 ± 1.9	—	966 ± 47	332 ± 16	59.3 ± 4.5	47.4 ± 4.4
	M	14.4 ± 3.4	—	812 ± 76	279 ± 25	48.7 ± 2.4	38.8 ± 3.2
significance of inoculation (I)		**		*	***	**	***
arsenic (As)		***		***	NS	***	***
I × As		NS		NS	NS	**	NS

^a NM and M represent nonmycorrhizal and mycorrhizal treatments, respectively. ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$; NS, not significant. ^b —, not detected.

inoculation decreased AR activities in plants. Further confirmation for such observation and elucidation of the influence mechanisms are necessary.

Suppressed reduction of As(V) could be significantly important for mycorrhizal plants to achieve As resistance. The toxic effect of As is highly dependent on its chemical speciation. Arsenite is generally believed to be more toxic to organisms than As(V) (1, 5). Moreover, conversion of As(V) to As(III) can generate reactive oxygen species (ROS) and result in damage to DNA, proteins, and lipids (2). Antioxidative enzymes are considered as an important defense system to avoid the damages to the ROS (26). We observed in this study that mycorrhizal inoculation markedly decreased POD and SOD activities in maize roots, which is consistent with the results of lower As(III) concentration and AR activities in roots. It can be expected that suppressed reduction of As(V) to As(III) in mycorrhizal maize reduced the generation of ROS, and therefore, less antioxidative enzymes were synthesized. However, our results showed that mycorrhizal inoculation increased P concentration in both roots and shoots (Table 1). It is possible that a high plant-P status protects plant membranes from As-induced oxidative stress (27), resulting in the low SOD and POD activities as evidenced by the results of the present study (Table 4).

Previous studies have confirmed that As can be metabolized from inorganic to organic forms by a wide range of bacteria, fungi, yeasts, and algae (1, 28). In addition, methylation of As has been observed in terrestrial plants (26). In this research, organic As species were detected in the plant and soil samples after plant harvest. Concentrations of DMA were found to be lower in the soils and roots for mycorrhizal than nonmycorrhizal treatment. MMA was only detected for nonmycorrhizal treatment. These results suggest that mycorrhizal inoculation might inhibit the methylation of inorganic As or enhance the demethylation of organic As. AM fungi can exude substances and result in a selective effect on the microbial community in rhizosphere soil and modify the microbial communities (29). Such processes might influence the amount and activity of soil microorganisms related to As methylation and demethylation. It has also been observed that methylation of As(V) by plants can take place under P-deficient conditions (26, 28). A higher P concentration and a lower organic As concentration were found in mycorrhizal roots of maize in this experiment; however, the mechanisms involved needs further investigation.

To our knowledge, this study is the first to report the effects of mycorrhizal inoculation on As speciation in plants. Mycorrhizal inoculation changed the uptake and accumulation of As(V) and As(III) in maize. More As(V) and less As(III) were accumulated in mycorrhizal roots compared to those in nonmycorrhizal roots, and shoot concentrations of As(V) and As(III) were all obviously lower for mycorrhizal compared to those for nonmycorrhizal plants. The findings of this study have contributed to our understanding of the role of AM fungi in As resistance in plants. Inhibited uptake of As(V) and reduction of As(V) to As(III) reduced the accumulation of As(III), the most toxic As species, in mycorrhizal plants. Mycorrhizal inoculation subsequently decreased the oxidation stress generated from As(V) reduction and alleviated As toxicity to the host plant.

LITERATURE CITED

- (1) Fitz, W. J.; Wenzel, W. W. Arsenic Transformations in the soil-rhizosphere-plant system: fundamentals and potential application to phytoremediation. *J. Biotechnol.* **2002**, *99*, 259–278.
- (2) Meharg, A. A.; Hartley-Whitaker, J. Arsenic uptake and metabolism in arsenic resistant and nonresistant plant species. *New Phytol.* **2002**, *154*, 29–43.
- (3) Zavala, Y. J.; Gerads, R.; Gurleyuk, H.; Duxbury, J. M. Arsenic in rice: II. Arsenic speciation in USA grain and implications for human health. *Environ. Sci. Technol.* **2008**, *42*, 3861–3866.
- (4) Meharg, A. A.; Jardine, L. Arsenite transport into paddy rice (*Oryza sativa*) roots. *New Phytol.* **2003**, *157*, 39–44.
- (5) Chakrabarty, D.; Trivedi, P. K.; Misra, P.; Tiwari, M.; Shri, M.; Shukla, D.; Kumar, S.; Rai, A.; Pandey, A.; Nigam, D.; Tripathi, R. D.; Tuli, R. Comparative transcriptome analysis of arsenate and arsenite stresses in rice seedlings. *Chemosphere* **2009**, *74*, 688–702.
- (6) Smith, S. E.; Read, D. J. *Mycorrhizal Symbiosis*, 2nd ed.; Academic Press: London, U.K., 1997.
- (7) Gonzalez-Chavez, C.; Harris, P. J.; Dodd, J.; Meharg, A. A. Arbuscular mycorrhizal fungi confer enhanced arsenate resistance on *Holcus lanatus*. *New Phytol.* **2002**, *155*, 163–171.
- (8) Leung, H. M.; Ye, Z. H.; Wong, M. H. Interactions of mycorrhizal fungi with *Pteris vittata* (As hyperaccumulator) in As-contaminated soils. *Environ. Pollut.* **2006**, *139*, 1–8.
- (9) Trotta, A.; Falaschi, P.; Cornara, L.; Minganti, V.; Fusconi, A.; Drava, G.; Berta, G. Arbuscular mycorrhizae increase the arsenic translocation factor in the As hyperaccumulating fern *Pteris vittata* L. *Chemosphere* **2006**, *65*, 74–81.
- (10) Ahmed, F. R. S.; Killham, K.; Alexander, I. Influences of arbuscular mycorrhizal fungus *Glomus mosseae* on growth and nutrition of lentil irrigated with arsenic contaminated water. *Plant Soil* **2006**, *283*, 33–41.
- (11) Chen, B. D.; Xiao, X. Y.; Zhu, Y. G.; Smith, F. A.; Xie, Z. M.; Smith, S. E. The arbuscular mycorrhizal fungus *Glomus mosseae* gives contradictory effects on phosphorus and arsenic acquisition by *Medicago sativa* Linn. *Sci. Total Environ.* **2007**, *379*, 226–234.
- (12) Liu, Y.; Zhu, Y. G.; Chen, B. D.; Christie, P.; Li, X. L. Yield and arsenate uptake of arbuscular mycorrhizal tomato colonized by *Glomus mosseae* BEG167 in As spiked soil under glasshouse conditions. *Environ. Int.* **2005**, *31*, 867–873.
- (13) Ultra, V. U.; Tanaka, S.; Sakurai, K.; Iwasaki, K. Effects of arbuscular mycorrhiza and phosphorus application on arsenic toxicity in sunflower (*Helianthus annuus* L.) and on the transformation of arsenic in the rhizosphere. *Plant Soil* **2007**, *290*, 29–41.
- (14) Hartley-Whitaker, J.; Ainsworth, G.; Meharg, A. A. Copper and arsenate-induced oxidative stress in *Holcus lanatus* L. clones with differential sensitivity. *Plant Cell Environ.* **2001**, *24*, 713–722.
- (15) Srivastava, M.; Ma, L. Q.; Singh, N.; Singh, S. Antioxidant responses of hyper-accumulator and sensitive fern species to arsenic. *J. Exp. Bot.* **2005**, *56*, 1335–1342.
- (16) Mukhopadhyay, R.; Rosen, B. P. Arsenate reductases in prokaryotes and eukaryotes. *Environ. Health Persp.* **2002**, *110*, 745–748.
- (17) Duan, G. L.; Zhu, Y. G.; Tong, Y. P.; Cai, C.; Kneer, R. Characterization of arsenate reductase in the extract of roots and fronds of Chinese brake fern, an arsenic hyperaccumulator. *Plant Physiol.* **2005**, *138*, 461–469.
- (18) Liu, Y.; Wang, H. B.; Wong, M. H.; Ye, Z. H. The role of arsenate reductase and superoxide dismutase in As accumulation in four *Pteris* species. *Environ. Int.* **2009**, *35*, 491–495.
- (19) Phillips, J. M.; Hayman, D. S. Improved procedures for clearing and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. Br. Mycol. Soc.* **1970**, *55*, 158–161.
- (20) Lynch, J. M. *Soil Rhizosphere*; John Wiley and Sons: New York, 1990.
- (21) Zhang, W. H.; Cai, Y.; Tu, C.; Ma, L. Q. Arsenic speciation and distribution in an arsenic hyperaccumulating plant. *Sci. Total Environ.* **2002**, *300*, 167–177.

- (22) Polle, A.; Chakrabarti, K.; Schurmann, W.; Rennenberg, H. Composition and properties of hydrogen-peroxide decomposing systems in extracellular and total extracts from needles of Norway spruce (*Picea-Abies* L, Karst). *Plant Physiol.* **1990**, *94*, 312–319.
- (23) Pickering, I. J.; Prince, R. C.; George, M. J.; Smith, R. D.; George, G. N.; Salt, D. E. Reduction and coordination of arsenic in Indian mustard. *Plant Physiol.* **2000**, *122*, 1171–1177.
- (24) Lee, Y. J.; George, E. Contribution of mycorrhizal hyphae to the uptake of metal cations by cucumber plants at two levels of phosphorus supply. *Plant Soil* **2005**, *278*, 361–370.
- (25) Duan, G. L.; Zhou, Y.; Tong, Y. P.; Mukhopadhyay, R.; Rosen, B. P.; Zhu, Y. G. A CDC25 homologue from rice functions as an arsenate reductase. *New Phytol.* **2007**, *174*, 311–321.
- (26) Sneller, F. E. C.; Van Heerwaarden, L. M.; Kraaijeveld-Smit, F. J. L.; Ten Bookum, W. M.; Koevoets, P. L. M.; Schat, H.; Verkleij, J. A. C. Toxicity of arsenate in *Silene vulgaris*, accumulation and degradation of arsenate-induced phytochelatins. *New Phytol.* **1999**, *144*, 223–232.
- (27) Gunes, A.; Pilbeam, D.; Inal, A. Effect of arsenic-phosphorus interaction on arsenic-induced oxidative stress in chickpea plants. *Plant Soil* **2009**, *314*, 211–220.
- (28) Nissen, P.; Benson, A. A. Arsenic metabolism in freshwater and terrestrial plants. *Physiol. Plantarum* **1982**, *54*, 446–450.
- (29) Duponnois, R.; Colombet, A.; Hien, V.; Thioulouse, J. The mycorrhizal fungus *Glomus intraradices* and rock phosphate amendment influence plant growth and microbial activity in the rhizosphere of *Acacia holosericea*. *Soil Biol. Biochem.* **2005**, *37*, 1460–1468.

Received for Review January 9, 2009. Revised manuscript received February 26, 2009. Accepted February 26, 2009. This work was funded by the National Natural Science Foundation of China (Projects 40730740, 20621703) and the National Basic Research Program (2009CB421603).