Selective 6-O-Debenzylation of Mono- and Disaccharide Derivatives Using ZnCl_2 - Ac_2O -HOAc

Guangbin Yang, Xianglan Ding, and Fanzuo Kong*

Research Center for Eco-Environmental Sciences, Academia Sinica, P.O.Box 2871,

Beijing 100085, P. R. China

Abstract: Freshly fused ZnCl_2 in Ac_2O /HOAc at room temperature has been used for 6-O-debenzylation of mono- and disaccharide derivatives giving yields more than 80%. Notably, allyl, acetyl, benzoyl, tosyl, thiono groups are unaffected under the designated conditions.
© 1997 Elsevier Science Ltd.

Many important bioactive natural products contain $1 \rightarrow 6$ glycosidic linkages¹⁻⁶. Selective 6-O-debenzylation of methyl glycoside derivatives is a very useful procedure for the synthesis of $1 \rightarrow 6$ linked oligosaccharides since it can substantially simplify the reaction pathway. Although the acetolysis of benzyl ethers initialized by Lewis acid or proton is well known, for example, by $\text{FeCl}_3/\text{Ac}_2\text{O}$ ⁷, $\text{ZnI}_2/\text{Ac}_2\text{O}$ ⁸, and $\text{H}_2\text{SO}_4/\text{Ac}_2\text{O}$ ⁹, the selectivity and yield are not satisfactory. The direct 6-O-debenzylation using trimethylsilyl trifluoromethanesulfonate/acetic anhydride ($\text{TMSOTf}/\text{Ac}_2\text{O}$)¹⁰ at -40°C has been shown to be temperature dependent, and secondary benzyl ether linkages can also be cleaved as the temperature is raised. Iodotrimethylsilane is another catalyst for direct 6-O-debenzylation¹¹ giving moderate to good yield, but the secondary benzyl groups can also be cleaved as the reaction time is long. In our recent studies, we found that ZnCl_2 (7.5-10eq)- Ac_2O -HOAc was a highly efficient reagent for 6-O-debenzylation at room temperature. Compounds of diverse structure including methyl α - and β -D-glucopyranoside, methyl α -D-galacto, manno, talopyranoside and methyl $1 \rightarrow 6$ linked disaccharide were tried. Notably, allyl, acetyl, benzoyl, tosyl, thiono groups all survived under the reaction conditions, no anomerization and glycosidic bond breaking were observed (see Table I for results). Also, 6-O-debenzylation of monosaccharides having one free

hydroxy group (see **2** and **4**) was very successfully performed giving the products with two potential hydroxy groups capable of building branched oligosaccharides. It was also found that a large excess of freshly fused ZnCl_2 (20eq) and a long reaction time (4h) would simultaneously cleave the methyl glycosidic bond while the secondary benzyl groups were still not changed (see Table I for results). The typical procedure for selective 6-O-debenzylation is as follows: To a solution of 6-O-benzylated sugar (50—200mg) in 1 ml $\text{Ac}_2\text{O}/\text{HOAc}$ (2:1) was added a solution of freshly fused ZnCl_2 (7.5—10 eq) in 1 ml $\text{Ac}_2\text{O}/\text{HOAc}$ (2:1), the mixture was stirred at room temperature for 2h, TLC (3:1 petroleum ether-ethyl acetate) indicated that the reaction was complete. Water was added, and the mixture was extracted with dichloromethane three times, washed with saturated sodium carbonate, then water, dried with Na_2SO_4 , and concentrated to give a syrup. Purification of the syrup by column chromatography (3:1 petroleum ether—ethyl acetate) yielded the pure product (yield >80%). The structures of most of the starting materials and products were simple, among which known compounds **1**, **1**^{'12}, **2**¹³, **3**, **3**['], **3**^{'14}, **5**, **5**^{'15}, **5**^{'16}, **7**¹⁷, **9**, **9**^{'18}, and **10**, **10**^{'19} were identified by comparison of their physical data to the reported values in literature, while new compounds were characterized by optical rotations, elemental analyses, and ^1H NMR spectrometry²⁰. Compared to the starting materials, the products gave ^1H NMR spectra showing a downfield chemical shift of H-6 and upfield shift of the acetyl methyl signal. In summary here we report a very effective and facile method for selective 6-O-debenzylation and this method can be used in the synthesis of oligosaccharides.

Table I Results for selective 6-O-debenzylation using ZnCl_2 - Ac_2O - HOAc

Compound	1	2	3	4	5	6	7	8	9	10
ZnCl_2 (eq)	7.5	10	7.5	10	7.5	10	10	10	7.5	10
Product	1 [']	2 [']	3 [']	4 [']	5 [']	6 [']	7 [']	8 [']	9 [']	10 [']
Yield	85%	90%	80%	86%	82%	88%	92%	94%	94%	81%
ZnCl_2 (eq)		20	20		20			20		
Product		2 ^{''}	3 ^{''}		5 ^{''}			8 ^{''}		
Yield		86%	90%		87%			92%		
			$\alpha:\beta$ 4:1		$\alpha:\beta$ 6:1

1-10 R=Bn R'=H R''=Me

1'-10' R=Ac R'=Ac R''=Me

2'',3'',5'',8'' R=Ac R'=Ac R''=Ac

ACKNOWLEDGEMENT

Project 29672049 was supported by The National Natural Science Foundation of China.

REFERENCES AND NOTES

1. Staaf, M.; Widmalm, G.; Yang, Z. and Hufninen, E. *Carbohydr. Res.*, **1996**, 291, 155.
2. Jain, R.K. and Matta, K.L. *Carbohydr. Res.*, **1996**, 282, 101.
3. Riccio, R.; Kinnel, R.B.; Biofulco, G. and Scheuer, P.J. *Tetrahedron lett.*, **1996**, 37, 1979.
4. Nakahara, Y.; Shibayama, S.; Nakahara, Y. and Ogawa, T. *Carbohydr. Res.*, **1996**, 280, 67.
5. Nicolaou, K.C.; Winssinger, N.; Pastor, J. and Deroose, F. *J. Am. Chem. Soc.*, **1997**, 119, 449.
6. Wyss, D.F.; Choi, J.S. and Wagner, G. *Science*, **1995**, 269, 1273.
7. Ganem, B. and Small, V.R.Jr. *J. Org. Chem.*, **1974**, 39, 3728.
8. Edith, S. and Burton, G. *J. Chem. Res. Synop.*, **1990**, 8, 248.
9. Sakai, J.I.; Takeda, T. and Ogihara, Y. *Carbohydr. Res.*, **1981**, 95, 125.

10. Angibeaud, P. and Utile, J.P. *Synthesis*, **1991**, 737.
11. Klemer, A.; Bieber, M. and Wilburs, H. *Liebigs Ann. Chem.*, **1983**, 1416
12. Shah, R.; Baptista, J. and Krepinsky, J.J. *J. Carbohydr. Chem.*, **1987**, 6(4), 645.
13. Liu, J.; Kong, F. and Cao, L. *Carbohydr. Res.*, **1993**, 240, 295.
14. Nashed, M.A. and Anderson, L. *Carbohydr. Res.*, **1976**, 51(1), 65.
15. Koto, S.; Morishima, N. and Zen, S. *Bull. Chem. Soc. Jpn.*, **1982**, 55(4), 1092.
16. Frechet, J.M. and Schuerch, C. *J. Am. Chem. Soc.*, **1972**, 94(2), 604.
17. Kong, F. and Su, B. *Carbohydr. Res.*, **1985**, 142(1), 152.
18. Koto, S.; Hamada, Y. and Zen, S. *Chem. Lett.*, **1975**, 6, 587.
19. Zhu, X.; Ding, P. and Cai, M. *Tetrahedron:Asym.*, **1996**, 7(10), 2833.
20. All new compounds gave satisfactory elemental analysis results. Selected ^1H NMR (CDCl_3 , Me_4Si as internal standard) data are as follows: **2'**. 7.36-7.24(m, 10H, Ph-H), 5.34(d, 1H, $J_{2,3}$ 1.7 Hz, H-2), 4.94, 4.75(AB_q, 2H, 2J 11.7 Hz, PhCH_2), 4.80(s, 1H, H-1), 4.75, 4.67(AB_q, 2H, 2J 11.7 Hz, PhCH_2), 4.40-4.18(m, 2H, H-6), 3.92-3.78(m, 3H, H-3,4,5), 3.32(s, 3H OCH_3), 2.10, 2.00(2s, 6H, 2COCH_3). **2''**. 7.38-7.24(m, 10H, Ph-H), 6.18(d, 1H, $J_{1,2}$ 0.9 Hz, H-1), 5.30(dd, 1H, $J_{2,3}$ 1.8 Hz, H-2), 4.92, 4.69(AB_q, 2H, 2J 12.0 Hz, PhCH_2), 4.76, 4.56(AB_q, 2H, 2J 12.0 Hz, PhCH_2), 4.32-3.32(m, 3H, H-3,4,5), 2.10, 2.08, 2.00(3s, 9H, 3COCH_3). **4'**. 8.04(d, 2H, J 8.9 Hz, Ph-H of Bz), 7.62-7.12(m, 8H, Ph-H), 5.85(t, 1H, $J_{3,4}$ 9.4 Hz, H-3), 5.05(dd, 1H, $J_{2,3}$ 9.4 Hz, H-2), 4.91(d, 1H, $J_{1,2}$ 3.7 Hz, H-1), 4.56, 4.48(AB_q, 2H, 2J 11.7 Hz, PhCH_2), 4.36(dd, 1H, $J_{6,6'}$ 12.5 Hz, $J_{5,6}$ 2.5 Hz, H-6), 4.20(dd, 1H, $J_{5,6'}$ 4.2 Hz, H-6'), 4.04-3.95(m, 1H, H-5), 3.76(t, 1H, $J_{4,5}$ 9.4 Hz, H-4), 3.43(s, 3H, OCH_3), 2.10, 1.96(2s, 6H, 2COCH_3). **6'**. 7.38-7.30(m, 5H, Ph-H), 5.08(t, $J_{3,4}$ 8.4 Hz, H-3), 5.06(s, 1H, H-1), 4.86, 4.60(AB_q, 2H, 2J 11.4 Hz, PhCH_2), 4.76(d, 1H, $J_{2,3}$ 7.6 Hz, H-2), 4.34(dd, 1H, $J_{6,6'}$ 12.3 Hz, $J_{5,6}$ 1.9 Hz, H-6), 4.22(dd, 1H, $J_{5,6'}$ 4.8 Hz, H-6'), 3.88-3.79(m, 1H, H-5), 3.59(dd, 1H, $J_{4,5}$ 9.5 Hz, H-4), 3.40(s, 3H, OCH_3), 2.00(s, 3H, COCH_3). **7'**. 7.72(d, 2H, J 8.7 Hz, Ph-H of Ts), 7.52-6.81(m, 12H, Ph-H), 4.85(dd, 1H, $J_{2,3}$ 3.4 Hz, $J_{3,4}$ 8.9 Hz, H-3), 4.70-4.47(m, 5H, 2PhCH_2 , H-1), 4.40-3.60(m, 5H, H-2,4,5,6), 3.32(s, 3H, OCH_3), 2.38(s, 3H, CH_3), 2.03(s, 3H, COCH_3). **8'**. 8.10(d, 2H, J 6.8 Hz, Ph-H of Bz), 7.60-7.28(m, 8H, Ph-H), 5.85-5.75(m, 1H, $\text{CH}_2=\text{CH}$), 5.50(dd, 1H, $J_{2,3}$ 3.3 Hz, H-3), 5.22(m, 1H, J_{trans} 15.0 Hz, 2J 1.3 Hz, $\text{HCH}=\text{CH}$), 5.09(m, 1H, J_{cis} 8.3 Hz, $\text{HCH}=\text{CH}$), 4.76(d, 1H, $J_{1,2}$ 0.5 Hz, H-1), 4.73, 4.56(AB_q, 2H, 2J 9.2 Hz, PhCH_2), 4.35(d, 2H, $J_{5,6}$ 2.7 Hz, H-6), 4.05-4.12(m, 2H, $\text{CH}_2=\text{CH}-\text{CH}_2$), 4.05(t, 1H, $J_{3,4}$ $J_{4,5}$ 8.7 Hz, H-4), 3.92(dd, 1H, $J_{2,3}$ 2.0 Hz, H-2), 3.40(s, 3H, OCH_3), 2.12(s, 3H, COCH_3). **8''**. 8.11(d, 2H, J 8.4 Hz, Ph-H of Bz), 7.66-7.12(m, 8H, Ph-H), 6.15(d, 1H, $J_{1,2}$ 1.8 Hz, H-1), 5.85-5.75(m, 1H, $\text{CH}_2=\text{CH}$), 5.51(dd, $J_{2,3}$ 4.2 Hz, H-3), 5.20(m, 1H, J_{trans} 16.3 Hz, 2J 0.9 Hz, $\text{HCH}=\text{CH}$), 5.10(m, 1H, J_{cis} 9.8 Hz, $\text{HCH}=\text{CH}$), 4.74, 4.60(AB_q, 2H, 2J 11.6 Hz, PhCH_2), 4.12(t, 1H, $J_{3,4}$ $J_{4,5}$ 9.3 Hz, H-4), 4.12-4.00(m, 3H, H-5, $\text{CH}_2=\text{CH}-\text{CH}_2$), 3.91(dd, 1H, $J_{2,3}$ 0.6 Hz, H-2), 2.14, 2.08(2s, 6H, COCH_3).

(Received in China 15 April 1997; revised 29 May 1997; accepted 1 July 1997)