Mercury in leaf litter in typical suburban and urban broadleaf forests in China

Zhenchuan Niu, Xiaoshan Zhang, Zhangwei Wang*, Zhijia Ci

Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China. E-mail: caoren03@yahoo.com.cn

Received 30 January 2011; revised 11 April 2011; accepted 14 April 2011

Abstract

To study the role of leaf litter in the mercury (Hg) cycle in suburban broadleaf forests and the distribution of Hg in urban forests, we collected leaf litter and soil from suburban evergreen and deciduous broadleaf forests and from urban forests in Beijing. The Hg concentrations in leaf litter from the suburban forests varied from 8.3 to 205.0 ng/g, with an average (avg) of (49.7 ± 36.9) ng/g. The average Hg concentration in evergreen broadleaf forest leaf litter (50.8 ± 39.4) ng/g was higher than that in deciduous broadleaf forest leaf litter (25.8 ± 10.1) ng/g. The estimated Hg fluxes of leaf litter in suburban evergreen and deciduous broadleaf forests were 179.0 and 83.7 mg/(ha·yr), respectively. The Hg concentration in organic horizons (O horizons) (263.1 ± 237.2) ng/g was higher than that in eluvial horizons (A horizons) (83.9 ± 52.0) ng/g. These results indicated that leaf litterfall plays an important role in transporting atmospheric mercury to soil in suburban forests. For urban forests in Beijing, the Hg concentrations in leaf litter ranged from 8.8–119.0 (avg 28.1 ± 16.6) ng/g, with higher concentrations at urban sites than at suburban sites for each tree. The Hg concentrations in surface soil in Beijing were 32.0–25300.0 ng/g and increased from suburban sites to urban sites, with the highest value from Jingshan (JS) Park at the centre of Beijing. Therefore, the distribution of Hg in Beijing urban forests appeared to be strongly influenced by anthropogenic activities.

Key words: mercury; broadleaf forests; leaf litter; soil; flux

DOI: 10.1016/S1001-0742(10)60669-9

Citation: Niu Z C, Zhang X S, Wang Z W, Ci Z J, 2011. Mercury in leaf litter in the typical suburban and urban broadleaf forests, China. 23(12): 2042–2048

Introduction

Mercury (Hg) is a global contaminant, likely spreading across the world through long-range atmospheric transport. Every year, thousands of tons of Hg are emitted into the atmosphere from natural and anthropogenic sources. Atmospheric Hg may be deposited into a forest canopy in gaseous and aerosol forms, and deposited Hg⁰ and reactive gaseous Hg (RGM) may be taken up by leaf stomata (Lindberg et al., 1979). Current research has identified that the strength of known atmospheric Hg sources is greater than previous estimations (Gustin et al., 2000; Engle et al., 2001; Pirroni et al., 2001), and vegetation is regarded as the missing sink in global Hg mass balance (Gustin et al., 2008). It is estimated that aboveground vegetation sequesters over 1000 tons Hg from the atmosphere every year (Obrist, 2007).

Litterfall is a major component of both energy flow and nutrient recycling in forest ecosystems. Global flux of Hg by way of litterfall is estimated to be 2400–6000 tons/yr, and is suggested to be the largest Hg flux to forest floors (Lindberg et al., 2004). Leaves account for 50%–84% of litterfall biomass (Silva-Filho et al., 2006) and their Hg

concentrations are the highest among litterfall components (Nóvoa-Muñoz et al., 2008). Consequently, leaves are a significant determinant factor of litterfall flux and studies about the role of leaf litter on the Hg cycle in forests are necessary. Researchers from countries such as the United States of America (Rea et al., 1996, 2002; Sheehan et al., 2006), Canada (St Louis et al., 2001; Graydon et al., 2008), Sweden (Iverfeldt, 1991; Munthe et al., 1995), Spain (Nóvoa-Muñoz et al., 2008), and Brazil (Roulet et al., 1999; Fostier et al., 2003; Silva-Filho et al., 2006), have studied the role of litterfall in transferring atmospheric Hg to soils in boreal, temperate, and tropical forests. In China, Wang et al. (2009) and Fu et al. (2010) reported the Hg concentrations and fluxes of litterfall in southwestern subtropical forests. However, regional and even nationwide investigations on Hg concentrations in leaf litter and soil in Chinese forests have not been conducted.

Mercury in surface soil is an indicator of soil quality, while Hg in foliages at urban sites is an indicator of air quality (Barghigiani et al., 1991; Zhang et al., 2006). In addition, mercury in surface soil and foliage is a potential source of atmospheric and soil Hg, respectively (Lindberg et al., 2004). During the past three decades, Beijing has been undergoing fast economic development and urban construction. Soil and atmospheric Hg pollution has been

* Corresponding authors. E-mail: wangzhw@rcees.ac.cn

studied in some urban areas (Liu et al., 2002; Wang et al., 2006; Zhang et al., 2006; Chen et al., 2010). It is also necessary, however, to study the distribution of Hg in leaf litter and soil in urban forests. Therefore, the goals of this study were to: (1) investigate the Hg concentrations in leaf litter and soil of evergreen and deciduous broadleaf forests in the suburban mountains of Chinese cities; and (2) investigate Hg distribution in leaf litter and surface soil within different forms of land use in Beijing urban forests.

1 Materials and methods

1.1 Site location

The study sites were divided into suburban and urban forests according to the influential extent of human activities. Suburban forests included both evergreen and deciduous broadleaf forests. Twenty-two sites were located at the suburban mountainous area of 19 cities in 8 provinces in China. Urban forest sites in Beijing were situated in the vicinity of landfills (outside the 5th Ring Road), universities, parks and streets, which contained 4–5 sites for each type of land use. The abbreviations and locations of these sites are presented in Table 1 and Fig. 1.

1.2 Sample collection

Samples of leaf litter and soils in eluvial horizons (A horizons) were collected from every suburban forest, and samples in organic horizons (O horizons) were also collected from some forests. The main tree species in the suburban evergreen broadleaf forests include *Castanopsis carlesii* (Hemsl.) Hayata, *Schima superba* Gardn. et Champ, *Castanopsis fargesii* Franch, *Cyclobalanopsis myrsinaefolia* (Bl.) Oerst, *Lithocarpus glaber* (Thunb.) Nakai, *Cinnamomum camphora* (Linn.) Presl and *Tsoongiodendron odoratum* Chun. In deciduous broadleaf forests, the main tree species were *Fraxinus mandschurica* Rupr.,

Populus davidiana Dode, *Betula platyphylla* Suk., *Ulmus pumila* L., *Quercus liaotungensis* Koidz. and *Acer truncatum* Bunge.

In Beijing urban forests, broadleaf trees *Sophora japonica* L., *Populus tomentosa* Carr., *Ginkgo biloba* L. were selected. For comparison, coniferous trees *Sabina chinensis* L. and *Pinus tabulaeformis* Carr. were also selected.

Fig. 1 Location of suburban forest sites in China and urban forest sites in Beijing. Abbreviations of sites are presented in Table 1.

Table 1 Abbreviations of suburban forest sites in China and urban forest sites in Beijing

Suburban forests in China				Urban forests in Beijing	
Forest type	Site	City/County	Province	Land use	Site
Evergreen broadleaf	Tieshanping (TSP)	Chongqing	Chongqing	Landfill	Gaoantun (GAT)
	Daling Mountain (DLM)	Dongguan	Guangdong		Beishenshu (BSS)
	Yinping Mountain (YPM)	Dongguan	Guangdong		Yonghezhuang (YHZ)
	Sanling Mountain (SLM)	Zhanjiang	Guangdong		Liulitun (LLT)
	Dinghu Mountain (DHM)	Zhaoqing	Guangdong	University	Tsinghua University (THU)
	Luchongguan (LCG)	Guiyang	Guizhou		Capital University of Economics and Business (CEB)
	Leigong Mountain (LGM)	Leishan	Guizhou		
	Maweizao Reservoir (MWZR)	Liuyang	Hunan		
	Daowu Mountain (DWM)	Liuyang	Hunan	Park	Beijing University of Technology (BUT)
	Heng Mountain (HM)	Hengyang	Hunan		Beijing Normal University (BNU)
	Wangxianling (WXL)	Chenzhou	Hunan		
	Hongqi Reservoir (HQR)	Ganzhou	Jiangxi		
	Wangzhai Reservoir (WZR)	Shangrao	Jiangxi	Street	Olympic Forest (OF)
	Xitianmu Mountain (XTMM)	Hangzhou	Zhejiang		Yuyuantan (YYT)
	Shuanglongdong (SLD)	Jinhua	Zhejiang		ChaoYang (CY)
	Fengyang Mountain (FYM)	Longquan	Zhejiang		Temple of Heaven (TH)
	Longwantan (LWT)	Yongjia	Zhejiang		Jing Shan (JS)
Deciduous broadleaf	Changbai Mountain (CBM)	Antu	Jilin		Zhongguan Cun (ZGC)
	Mang Mountain (MM)	Changping	Beijing		Beigao (BG)
	Badaling Greatwall (BDLG)	Yanqing	Beijing		Shuangqing (SQ)
	Xisancun (XSC)	Huairou	Beijing		Tiananmen (TAM)
	Hongluosi (HLS)	Huairou	Beijing		

Leaf samples were collected from five parallel individuals at each site. Surface soil (1–2 cm) was collected at each site from all four directions (north, south, west, and east) with a stainless steel shovel, and then was mixed together as the sample of one site.

Samples of leaf litter and soil were put in plastic zip-lock bags and transported to the laboratory immediately. Leaf litter was first rinsed by tap water, and then rinsed by ultra pure deionized water in laboratory. Leaf litter samples were dried at 55°C, ground, and then stored at –4°C for analysis. Soil samples were air-dried, ground, sieved and then stored at –4°C for analysis.

1.3 Sample analysis

The Hg concentrations in leaf litter and soil samples were measured by an RA-915⁺ Hg analyzer attached with PYRO-91 thermal decomposition accessory (Lumex Inc., Russia). The samples were thermally decomposed in an atomizer chamber at 750°C with aided catalytic action, and then the RA-915⁺ analyzer detected the Hg⁰. Each sample was analyzed three times and then averaged. In this study, instrument calibration curves covering the appropriate concentrations were confirmed by soil standards (GBW07404, 590 ng/g) and peach foliage standards (GBW08501, 40 ng/g), and control standard samples were checked every ten samples. The instrument detection limit was 5 pg for solid samples. The precisions, obtained from ten replicated determinations of standards, were 3.47% for soil and 2.83% for peach foliage.

1.4 Data analysis

Analysis of variance (ANOVA) was performed using SPSS 11.5 for windows. Data were presented as mean and standard deviation, and statistical significance was considered at $p < 0.05$.

2 Results and discussion

2.1 Hg concentrations in leaf litter and soil in Chinese suburban forests

The average Hg concentration in evergreen broadleaf forest leaf litter (50.8 ± 39.4 ng/g) was about two times the concentration in deciduous broadleaf forest litter (25.8 ± 10.1 ng/g). This may be the result of a longer growth period for evergreen plants, with previous studies indicating that Hg concentrations in foliage increased with time (Eriksen et al., 2003; Millhollen et al., 2006). The Hg concentration at SLD (119.9 ± 30.3 ng/g) was the highest among all the sites, with LCG (91.8 ± 78.2 ng/g) presenting the second highest concentration (Table 2). In this study, the Hg concentrations in leaf litter in Chinese suburban forests were 8.3–205.0 (avg 49.7 ± 36.9) ng/g, within the range of 8–250 ng/g reported in the United States of America, Canada, Brazil, Sweden, and French Guiana (Table 3), but lower than those reported in Chinese southwest forests (Wang et al., 2009).

The soil Hg concentrations in A horizons and O horizons in Chinese suburban forests were 23.7–181.7 (avg 83.9 ± 52.0) ng/g and 95.3–860.7 (avg 263.1 ± 237.2) ng/g, respectively. Most soil Hg concentrations in the A horizons were lower than the Natural Standard Value for Soil Environment Quality I of Hg (150 ng/g, GB15618–1995) and concentrations in 11 sites were lower than the background soil Hg value (65 ng/g) in China (SEPAC, 1990), which indicated that most A horizon soils in the suburban forests were not polluted by Hg. The concentrations in the O horizons were 1.2–19.1 times those in the A horizons. Nóvoa-Muñoz et al. (2008) also reported Hg concentrations in O horizons (216.9–248.2 ng/g) higher than those in A horizons (108.1–195.0 ng/g) in Spain. Wang et al. (2009) reported that soil Hg concentrations decreased with soil depth in Chinese southwest forests. Larssen et al. (2008) reported the organic horizon and the

Table 2 Hg concentrations in leaf litter and soil at each suburban forest site

Evergreen broadleaf forest				Deciduous broadleaf forest			
Site	Leaf litter (ng/g) mean $\pm$ SD (range)	Soil (ng/g)		Site	Leaf litter (ng/g) mean $\pm$ SD (range)	Soil (ng/g)	
		O horizon	A horizon			O horizon	A horizon
DLM	22.0 $\pm$ 12.7 (13.0–31.0)	–	31.0 $\pm$ 4.2	CBM	27.0 $\pm$ 8.8 (21.5–40.0)	860.7 $\pm$ 24.2	45.0 $\pm$ 1.4
DWM	45.5 $\pm$ 18.3 (23.0–68.0)	221.0 $\pm$ 7.0	181.7 $\pm$ 9.0	MM	21.3 $\pm$ 8.6 (13.3–33.0)	–	63.0 $\pm$ 1.4
DHM	46.6 $\pm$ 10.8 (34.0–64.0)	–	170.5 $\pm$ 2.1	BDLG	31.7 $\pm$ 25.0 (14.0–49.3)	–	40.0 $\pm$ 2.1
FYM	30.7 $\pm$ 7.5 (22.0–35.0)	99.3 $\pm$ 5.5	66.5 $\pm$ 3.5	XSC	27.4 $\pm$ 9.0 (15.0–36.3)	–	32.0 $\pm$ 1.4
HM	68.8 $\pm$ 41.4 (32.0–125.0)	130.7 $\pm$ 0.6	95.7 $\pm$ 3.1	HLS	25.0 $\pm$ 9.3 (15.7–39.0)	–	37.3 $\pm$ 2.1
HQR	25.0 $\pm$ 10.4 (18.0–37.0)	–	58.5 $\pm$ 3.5				
LGM	69.6 $\pm$ 55.5 (25.0–187.5)	323.3 $\pm$ 17.6	177.7 $\pm$ 11.7				
LWT	57.4 $\pm$ 15.6 (31.0–69.0)	–	61.0 $\pm$ 2.6				
LCG	91.8 $\pm$ 78.2 (34.0–205.0)	283.3 $\pm$ 7.8	121.3 $\pm$ 8.4				
MWZR	57.0 $\pm$ 13.8 (41.0–69.0)	95.3 $\pm$ 1.5	23.7 $\pm$ 1.5				
SLM	31.5 $\pm$ 21.9 (16.0–47.0)	–	39.5 $\pm$ 4.9				
SLD	119.9 $\pm$ 30.3 (91.5–147.0)	–	71.5 $\pm$ 2.1				
TSP	41.3 $\pm$ 12.4 (29.0–57.0)	–	79.0 $\pm$ 1.4				
WXL	67.5 $\pm$ 21.1 (41.0–86.0)	182.3 $\pm$ 4.2	155.0 $\pm$ 2.0				
WZR	21.1 $\pm$ 13.1 (8.3–47.0)	–	150.5 $\pm$ 0.7				
XTMM	43.0 $\pm$ 49.2 (9.7–151.0)	172.3 $\pm$ 3.2	94.3 $\pm$ 0.6				
YPM	16.9 $\pm$ 5.9 (9.4–22.0)	–	51.5 $\pm$ 2.1				

– indicates no samples.

Table 3 Comparison of Hg concentrations and fluxes in different countries

Country	Hg concentration (ng/g)	Hg flux (mg/(ha·yr))	Reference
China	leaf litter: 8.3–205, 49.7 ± 36.9 litter: 104.8 ± 18.6–135.1 ± 31.7	leaf litter: 179.0, 83.7	This study Wang et al., 2009
America	green foliage: 34.2 ± 7.2, litter: 53.2 ± 11.4 litter: 36 ± 8	litter: 130 litter: 158 ± 19, 114 ± 28	Rea et al., 1996 Rea et al., 2002
Canada	litter: 31.6–58.8 litter: 33–79	litter: 100, 101 litter: 110–220	Sheehan et al., 2006 St. Louis et al., 2001
Sweden	litter: 28.9–69.2 litter: 97.4–140.6, 20–80	litter: 86–105 litter: 250	Graydon et al., 2008 Iverfeldt, 1991
Brazil	litter: 33–140 green foliage: 40–152, leaf litter: 56–140 litter: 72–100		Munthe et al., 1995 Roulet et al., 1999
French Guiana	litter: 20–244 leaves: 32.4–114	litter: 1220 litter: 450 ± 100	Fostier et al., 2003 Silva-Filho et al., 2006 Mélières et al., 2003

illuviation horizon have clearly higher Hg concentrations than do other horizons in mineral soil in Norway. However, Roulet et al. (1999) reported that the rapid turnover of organic matter at the surface of tropical soil in the Brazilian Amazonian ecosystems limited the accumulation of Hg in the organic horizon, which was much lower than that in the underlying mineral horizons. The present study showed that the Hg concentration in the O horizon at CBM (860.7 ng/g) was much higher than that in the other evergreen broadleaf forests. Slow decomposition of litter due to low temperatures at CBM led the Hg in litterfall accumulated in the O horizon. Guo et al. (2006) reported that 95% of leaf and twig decomposition takes about 4.5–8.0 and 7.8–29.3 years at CBM, respectively.

2.2 Distribution of Hg in leaf litter and surface soil in Beijing forests

The Hg concentrations in leaf litter in Beijing were 28.1 ± 16.6 (8.8–119.0) ng/g, nearly the same as those in suburban deciduous broadleaf forests (25.8 ± 10.1) ng/g), but lower than those in suburban evergreen broadleaf forests (50.8 ± 39.4) ng/g). This may ascribe that the tree species in Beijing are mainly deciduous, and their leaf growth periods are shorter than those of evergreen plants. The Hg concentrations in leaf litter did not correlate

significantly with soil Hg concentrations for each species ($p > 0.05$), so Hg in leaves may originate mainly from the atmosphere. For *S. japonica*, the Hg concentrations at the streets, gardens, and universities were higher than those near landfills and suburban areas. The same results were found for *P. tomentosa*, *G. biloba*, *S. chinensis*, and *P. tabulaeformis* (Table 4). De Temmerman et al. (2007, 2009) showed that foliar Hg concentrations linearly correlate with atmospheric Hg concentrations, while Liu et al. (2002) showed that atmospheric Hg concentrations at the centre of Beijing ($7.9\text{--}34.9$ ng/m³) are higher than those in suburban ($5.3\text{--}12.4$ ng/m³) and rural ($2.5\text{--}5$ ng/m³) areas. Therefore, the difference of Hg concentrations in leaf litter for each species might roughly indicate the atmospheric Hg pollution among different land use forms.

At most sites, the Hg concentrations in *S. japonica*, *P. tomentosa* and *G. biloba* were higher than those in *S. chinensis* and *P. tabulaeformis*. It appeared, therefore, that the leaves of broadleaf trees accumulated more Hg than those of coniferous trees. This may result from the fewer stoma in the leaves of coniferous trees.

The Hg concentrations in surface soil in Beijing varied from 32.0 to 25300.0 ng/g, with a median of 129.5 ng/g and an average of 1502.1 ng/g. Li et al. (2010) reported that the Hg concentration in Beijing topsoil ranges from 12.1

Table 4 Hg concentrations in surface soils and leaf litter at different land use types in Beijing

Type	Site	Soil (ng/g)	Avg (ng/g)	Leaf litter (ng/g)									
				<i>S. japonica</i>	Avg	<i>P. tomentos</i>	Avg	<i>G. biloba</i>	Avg	<i>S. chinensis</i>	Avg	<i>P. tabulaeformis</i>	Avg
Suburban	MM	63.0 ± 0.4	44.1	33.0 ± 1.4	28.3	22.0 ± 1.4	29.2	—	—	13.3 ± 0.5	14.7	16.7 ± 0.5	21.4
	XSC	32.0 ± 1.4		30.7 ± 0.5		36.3 ± 0.9		—		15.0 ± 0.8		27.7 ± 0.9	
	HLS	37.3 ± 1.7		21.3 ± 2.1		29.3 ± 2.1		—		15.7 ± 0.5		19.7 ± 0.5	
Landfill	GAT	89.0 ± 5.4	99.4	28.7 ± 0.5	33.1	37.0 ± 0.0	28.0	11.7 ± 1.2	12.6	16.3 ± 0.9	14.8	13.7 ± 1.2	13.8
	LLT	94.3 ± 17.6		69.0 ± 1.6		26.0 ± 1.4		—		23.7 ± 1.2		21.0 ± 0.8	
	BSS	101.0 ± 12.6		22.3 ± 0.5		22.0 ± 0.8		13.7 ± 0.9		8.8 ± 0.8		9.4 ± 0.4	
University	YHZ	113.3 ± 11.8		12.3 ± 0.5		27.0 ± 1.6		12.3 ± 0.9		10.3 ± 0.5		11.0 ± 0.8	
	THU	255.0 ± 49.3	384.6	67.0 ± 2.2	44.8	42.3 ± 1.7	36.8	23.7 ± 1.2	18.8	21.0 ± 0.8	17.3	57.3 ± 0.5	29.6
	CEB	308.7 ± 30.0		40.0 ± 2.2		35.0 ± 0.8		15.3 ± 0.9		15.0 ± 0.8		19.0 ± 0.8	
Park	BUT	442 ± 55.7		32.0 ± 0.0		36.0 ± 0.8		15.7 ± 0.5		17.3 ± 0.5		19.0 ± 0.0	
	BNU	532.7 ± 49.3		40.0 ± 0.0		33.7 ± 0.5		20.7 ± 0.5		15.7 ± 0.5		23.0 ± 0.8	
	OF	117.0 ± 16.3	5420.2	49.3 ± 0.9	45.3	28.7 ± 1.2	37.6	11.3 ± 0.5	25.5	13.0 ± 0.8	16.7	52.3 ± 2.1	28.9
Street	YYT	137.3 ± 8.8		39.0 ± 0.0		44.0 ± 0.0		27.7 ± 1.2		13.0 ± 0.8		21.0 ± 1.4	
	CY	250.0 ± 43.3		37.0 ± 0.8		47.7 ± 1.2		23.3 ± 1.2		12.0 ± 0.8		16.7 ± 0.9	
	TH	1296.7 ± 9.4		41.0 ± 0.8		33.0 ± 0.8		27.7 ± 1.7		20.3 ± 0.5		22.0 ± 0.8	
Street	JS	25300.0 ± 408.2		60.0 ± 0.0		34.7 ± 0.5		37.3 ± 2.6		25.0 ± 1.4		32.3 ± 2.1	
	ZGC	81.3 ± 10.7	217.9	42.3 ± 1.2	46.3	32.3 ± 1.2	34.0	12.3 ± 0.5	18.7	13.3 ± 0.9	20.8	20.0 ± 1.4	45.0
	BG	121.7 ± 10.7		39.3 ± 0.5		30.0 ± 0.8		21.3 ± 1.2		23.0 ± 0.0		21.3 ± 0.5	
Street	SQ	262.0 ± 15.1		75.0 ± 1.6		38.0 ± 1.6		26.7 ± 0.5		34.0 ± 0.8		119.0 ± 0.8	
	TAM	406.7 ± 27.4		28.3 ± 1.2		35.7 ± 0.5		14.3 ± 0.5		12.7 ± 0.5		19.7 ± 0.5	

to 8487 ng/g, lower than the maximal value of 25300 ng/g in this study, and the median Hg concentration is 471.3 ng/g, much higher than the median concentration of 129.5 ng/g in this study. Most Hg concentrations were lower than the Natural Standard Value for Soil Environment Quality II of Hg (1000 ng/g, GB15618–1995), except the soil Hg concentrations in the TH and JS Parks. The median Hg concentration of 129.5 ng/g in surface soil was 61.9% higher than the reported background concentrations in Beijing (80 ng/g, Environmental Monitoring of China, 1994).

The Hg concentrations in surface soil increased from suburban sites to urban sites, with the highest value in the JS Park at the centre of Beijing. Soil Hg concentrations with different types of land use followed the sequence: classical park (TH, JS) > university > street > vicinity of landfill > suburban. These results indicated that the Hg distribution in Beijing was influenced by anthropogenic activities. Li et al. (2010) reported that the median Hg concentration in different regions occurred in the following order: urban center soil > suburban soil > rural soil, and thought that the regional Hg distribution was closely related to anthropogenic and industrial activities. The Hg concentrations in classical parks were relatively higher than those in other land use forms, and also higher than those in parks in many other countries (Table 5). This may relate to the use of gold amalgamation or cinnabar as paint in classical parks. Chen et al. (2010) also reported soil Hg concentrations in three classical parks (Beihai, Zhongshan, TH) in Beijing as 9400, 6500 and 4200 ng/g, respectively. In this study, the Hg concentrations in universities were relatively high, as reported by Chen et al. (2010) who inferred

that the use of Hg in medical and scientific apparatuses in cultural and educational areas was the main reason for this concentration. However, the Hg concentrations at the streets (except TAM) were not actually high. Chen et al. (2010) suggested that traffic was not a main source for Hg in roadside areas. The low Hg concentrations in the vicinity of landfills may ascribe to location, as all landfills were located outside the 5th Ring Road, far from urban areas.

2.3 Estimations of Hg fluxes of leaf litter in evergreen and deciduous broadleaf forests

In this study, the Hg flux of leaf litterfall was estimated by the biomass of leaf litter multiplied by the Hg concentration in leaf litter (Rea et al., 1996; Sheehan et al., 2006; Silva-Filho et al., 2006; Wang et al., 2009). Because of a lack of information on biomass of leaf litterfall in urban forests, we only estimated the fluxes in suburban evergreen and deciduous broadleaf forests. We averaged the data on biomass of leaf litter in evergreen and deciduous broadleaf forests reported by others, and obtained Hg fluxes of 179.0 and 83.7 mg/(ha·yr), respectively (Table 6). These fluxes were nearly at the same level of those reported in America, Canada and Sweden (Table 3), but lower than those reported in French Guiana (Mélières et al., 2003), and much lower than those reported in the Brazilian Amazon (Silva-Filho et al., 2006) (Table 3). The extremely high values at the two sites may be ascribed to the huge biomass of litterfall in tropical rainforests. The Hg fluxes of leaf litterfall in deciduous and evergreen broadleaf forests were nearly the same with a wet deposition flux of 84 mg/(ha·yr) and a dry deposition flux of 165 mg/(ha·yr) at CBM (Wan et al., 2009), respectively. Therefore, the input

Table 5 Soil Hg concentrations in urban forests at different countries

Site	Hg concentration (ng/g)	Reference
Beijing, China	32.0–25300, classical park: 1297–25300	This study
Beijing, China	16–966 (278)	Zhang et al., 2006
Beijing, China	22–9400, classical park: 1100–9400	Chen et al., 2010
Aveiro, Portugal	Park: 32–130 (59)	Rodrigues et al., 2006
Glasgow, England	Park: 310–5200 (1600)	Rodrigues et al., 2006
Ljubljana, Serbia	Park: 150–860 (410)	Rodrigues et al., 2006
Sevilla, Spain	Park: 110–1300 (420)	Rodrigues et al., 2006
Torino, Italy	Park: 210–900 (480)	Rodrigues et al., 2006
Uppsala, Sweden	Park: 15–1200 (350)	Rodrigues et al., 2006
Palermo, Italy	Park and greenland: 40–56000, media 1850	Manta et al., 2002

Table 6 Hg fluxes of leaf litter in evergreen and deciduous broadleaf forests

Broadleaf forest type	Biomass of leaf litter reported in China (kg/(ha·yr))	Average biomass (kg/(ha·yr))	Average Hg concentrations (ng/g)	Hg flux (mg/(ha·yr))	Reference
Evergreen	3764 2969.1 3033, 3499 4260 1876.18, 3970.88 3964 1788.07 2350 4031.4, 6143.2, 4155.4	3523.3	50.8	179.0	Hou et al., 1998 Lin et al., 1999 Yang et al., 2001 Guan et al., 2004 Chen et al., 2006 Wu, 2006 Fan et al., 2007 Wei et al., 2009 Pan et al., 2010
Deciduous	2549 4210, 3220, 3350 2890.8	3244.0	25.8	83.7	Zhang et al., 2008b Zhang et al., 2008a Liu et al., 2009

of Hg through leaf litterfall is important for the transport of atmospheric Hg to soil in Chinese forests.

The urban forest is an important forest type and is closely related with the daily life of humans. However, studies on litterfall in urban forests are limited, thus it is necessary to conduct research on the role of litterfall on the Hg cycle of urban forests.

3 Conclusions

In Chinese suburban forests, the Hg concentrations and fluxes in leaf litter in evergreen broadleaf forests were higher than those in deciduous broadleaf forests. The Hg concentrations in O horizons were higher than those in A horizons, and the Hg concentration in the O horizon at CBM was the highest among these sites. This data indicated that leaf litterfall is important for the transport of atmospheric Hg to soil in both evergreen and deciduous broadleaf forests.

The Hg concentrations in leaf litter at urban sites were higher than those at suburban sites for each plant. The Hg concentrations in surface soil in Beijing increased from suburban to urban areas, with the highest value at the center of Beijing. It therefore appears that the distribution of Hg in urban forests is strongly influenced by anthropogenic activities.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No. 40803033, 41073092). The authors would like to thank Yujing Mu, Yi Zhang, Zhanyi Wang and Bingwen Li from Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences for their suggestions and help in sample collection. The anonymous reviewers are acknowledged for their valuable comments.

References

- Barghigiani C, Ristori T, Bauleo R, 1991. Pinus as an atmospheric Hg biomonitor. *Environmental Technology*, 12(12): 1175–1181.
- Chen A L, Lin S Z, Chen S P, Chen Y Z, 2006. Comparison on litterfall character of *Phoebe bournei* and *Cunninghamia lanceolata* plantation in the field of successive planting *Cunninghamia lanceolata*. *Journal of Fujian College of Forestry*, 26(4): 289–293.
- Chen X, Xia X H, Wu S, Wang F, Guo X J, 2010. Mercury in urban soils with various types of land use in Beijing, China. *Environmental Pollution*, 158(1): 48–54.
- De Temmerman L, Claeys N, Roekens E, Guns M, 2007. Biomonitoring of airborne mercury with perennial ryegrass cultures. *Environmental Pollution*, 146(2): 458–462.
- De Temmerman L, Waegeneers N, Claeys N, Roekens E, 2009. Comparison of concentrations of mercury in ambient air to its accumulation by leafy vegetables: An important step in terrestrial food chain analysis. *Environmental Pollution*, 157(4): 1337–1341.
- Engle M A, Gustin M S, Zhang H, 2001. Quantifying natural source mercury emissions from the Ivanhoe Mining District, north-central Nevada, USA. *Atmospheric Environment*, 35(23): 3987–3997.
- Environmental Monitoring of China, 1994. The illustrated handbook of the background value of soil environment in the People's Republic of China. Chinese Environment Sciences Press, Beijing. 26–75.
- Ericksen J A, Gustin M S, Schorran D E, Johnson D W, Lindberg S E, Coleman J S, 2003. Accumulation of atmospheric mercury in forest foliage. *Atmospheric Environment*, 37(12): 1613–1622.
- Fan H B, Li Y Y, Su B Q, Lin D X, Liu C H, Jiang Z K, 2006. Allocation pattern of biomass and productivity in the mixed uneven-aged stands of Masson's pine and hardwood species. *Acta Ecologica Sinica*, 26(8): 2463–2473.
- Fostier A H, Cecon K, Forti M C, 2003. Urban influence on litterfall trace metals fluxes in the Atlantic Forest of São Paulo (Brazil). *Journal de Physique IV (Proceeding)*, 107(1): 491–494.
- Fu X W, Feng X B, Zhu W Z, Rothenberg S, Yao H, Zhang H, 2010. Elevated atmospheric deposition and dynamics of mercury in a remote upland forest of southwestern China. *Environmental Pollution*, 158(6): 2324–2333.
- Graydon J A, St Louis V L, Hintelmann H, Lindberg S E, Sandilands K A, Rudd J W M et al., 2008. Long-term wet and dry deposition of total and methyl mercury in the remote boreal ecoregion of Canada. *Environmental Science and Technology*, 42(22): 8345–8351.
- Guan L L, Zhou G Y, Zhang D Q, Liu J X, Zhang Q M, 2004. Twenty years of litter fall dynamics in subtropical evergreen broad-leaved forests at the Dinghushan forest ecosystem research station. *Acta Phytocologica Sinica*, 28(4): 449–456.
- Guo Z L, Zheng J P, Ma Y D, Li Q K, Yu G R, Han S J et al., 2006. Researches on litterfall decomposition rates and model simulating of main species in various forest vegetations of Changbai Mountains, China. *Acta Ecologica Sinica*, 26(4): 1037–1046.
- Gustin M S, Lindberg S E, Austin K, Coolbaugh M, Vette A, Zhang H, 2000. Assessing the contribution of natural sources to regional atmospheric mercury budgets. *Science of the Total Environment*, 259(1-3): 61–71.
- Gustin M S, Lindberg S E, Weisberg P J, 2008. An update on the natural sources and sinks of atmospheric mercury. *Applied Geochemistry*, 23(3): 482–493.
- Hou Y, Wang B S, Zhang H D, Li M G, 1998. The litterfall of the south subtropical evergreen broad-leaved forest in Heishiding nature reserve. *Ecologic Science*, 17(2): 14–18.
- Iverfeldt Å, 1991. Mercury in forest canopy throughfall water and its relation to atmospheric deposition. *Water, Air, and Soil Pollution*, 56(1): 553–564.
- Larssen T, de Wit H A, Wiker M, Halse K, 2008. Mercury budget of a small forested boreal catchment in southeast Norway. *Science of the Total Environment*, 404(2-3): 290–296.
- Li X H, Cheng H X, Zhao C D, Xu X B, 2010. Mercury contamination in the topsoil and subsoil of urban areas of Beijing, China. *Bulletin of Environmental Contamination and Toxicology*, 85(2): 224–228.
- Lin Y M, He J Y, Yang Z W, Liu C D, Lin P, Li Z J, 1999. The dynamics and production of litter falls of *Castanopsis eyrei* community in Wuyi Mountains. *Journal of Xiamen University (Natural Science)*, 38(2): 280–286.
- Lindberg S E, Jackson D R, Huckabee J W, Janzen S A, Levin M J, Lund J R, 1979. Atmospheric emission and plant uptake of mercury from agricultural soils near the Almaden

- Mercury Mine. *Journal of Environmental Quality*, 8(4): 572–578.
- Lindberg S E, Porcella D, Prestbo E, Friedli H, Radke L, 2004. The problem with mercury: too many sources, not enough sinks. *RMZ Materials and Geoenvironment*, 52: 1172–1176.
- Liu S L, Nadim F, Perkins C, Carley R J, Hoag G E, Lin Y H et al., 2002. Atmospheric mercury monitoring survey in Beijing, China. *Chemosphere*, 48(1): 97–107.
- Liu Y, Han S J, Lin L, 2009. Dynamic changes in soil nutrients of four types of forests in Changbai Mountains during litter decomposition. *Journal of Northeast Forestry University*, 37(8): 28–30.
- Manta D S, Angelone M, Bellanca A, Neri R, Sprovieri M, 2002. Heavy metals in urban soils: a case study from the city of Palermo (Sicily), Italy. *Science of the Total Environment*, 300(1-3): 229–243.
- Mélières M A, Pourchet M, Charles-Dominique P, Gaucher P, 2003. Mercury in canopy leaves of French Guiana in remote areas. *Science of the Total Environment*, 311(1-3): 261–267.
- Millhollen A G, Gustin M S, Obrist D, 2006. Foliar mercury accumulation and exchange for three tree species. *Environmental Science and Technology*, 40(19): 6001–6006.
- Munthe J, Hultberg H, Iverfeldt Å, 1995. Mechanisms of deposition of methylmercury and mercury to coniferous forests. *Water, Air, and Soil Pollution*, 80(1-4): 363–371.
- National Standards of the People's Republic of China, 1995. Environmental quality standard for soils. GB15618–1995.
- Nóvoa-Muñoz J C, Pontevedra-Pombal X, Martínez-Cortizas A, Gayoso E G R, 2008. Mercury accumulation in upland acid forest ecosystems nearby a coal-fired power-plant in Southwest Europe (Galicia, NW Spain). *Science of the Total Environment*, 394(2-3): 303–312.
- Obrist D, 2007. Atmospheric mercury pollution due to losses of terrestrial carbon pools? *Biogeochemistry*, 85(2): 119–123.
- Pan H, Huang S D, Hong W, Zhao K, Zhang Z H, 2010. Litterfalls amounts and seasonal dynamics of carbon returns in three Acacia plantations. *Journal of Fujian College of Forestry*, 30(2): 104–108.
- Pirroni N, Costa P, Pacyna J M, Ferrara R, 2001. Mercury emissions to the atmosphere from natural and anthropogenic sources in the Mediterranean region. *Atmospheric Environment*, 35(17): 2997–3006.
- Rea A W, Keeler G J, Scherbatskoy T, 1996. The deposition of mercury in throughfall and litterfall in the lake champlain watershed: A short-term study. *Atmospheric Environment*, 30(19): 3257–3263.
- Rea A W, Lindberg S E, Scherbatskoy T, Keeler G J, 2002. Mercury accumulation in foliage over time in two northern mixed-hardwood forests. *Water, Air, and Soil Pollution*, 133(1-4): 49–67.
- Rodrigues S, Pereira M E, Duarte A C, Ajmone-Marsan F, Davidson C M, Grčman H et al., 2006. Mercury in urban soils: A comparison of local spatial variability in six European cities. *Science of the Total Environment*, 368(2-3): 926–936.
- Roulet M, Lucotte M, Farella N, Serique G, Coelho H, Souza Passos C J et al., 1999. Effects of recent human colonization on the presence of mercury in Amazonian ecosystems. *Water, Air and Soil Pollution*, 112(3-4): 297–313.
- Sheehan K D, Fernandez I J, Kahl J S, Amirbahman A, 2006. Litterfall mercury in two forested watersheds at Acadia National Park, Maine, USA. *Water, Air, and Soil Pollution*, 170(1-4): 249–265.
- Silva-Filho E V, Machado W, Oliveira R R, Sella S M, Lacerda L D, 2006. Mercury deposition through litterfall in an Atlantic Forest at Ilha Grande, Southeast Brazil. *Chemosphere*, 65(11): 2477–2484.
- State Environment Protection Administration of China (SEPAC), 1990. The Background Levels of Element in soil in China. Chinese Environmental Science Press, Beijing.
- St Louis V L, Rudd J W M, Kelly C A, Hall B D, Rolfs K R, Scott K J et al., 2001. Importance of the forest canopy to fluxes of methyl mercury and total mercury to boreal ecosystems. *Environmental Science and Technology*, 35(15): 3089–3098.
- Wan Q, Feng X B, Lu J L, Zheng W, Song X J, Li P et al., 2009. Atmospheric mercury in Changbai Mountain area, northeastern China II. The distribution of reactive gaseous mercury and particulate mercury and mercury deposition fluxes. *Environmental Research*, 109(6): 721–727.
- Wang S X, Liu M, Jiang J K, Hao J M, Wu Y, Street D G, 2006. Estimate the mercury emissions from non-coal sources in China. *Chinese Journal of Environmental Science*, 27(12): 2401–2406.
- Wang Z W, Zhang X S, Xiao J S, Ci Z J, Yu P Z, 2009. Mercury fluxes and pools in three subtropical forested catchments, southwest China. *Environmental Pollution*, 157(3): 801–808.
- Wei L M, Yu D L, Chen Z R, 2009. Study on the dynamic variation of litter falls in Maolan Karst forest. *Journal of Nanjing Forestry University (Natural Science Edition)*, 33(3): 31–34.
- Wu Z X, 2006. Amount, component, and seasonal pattern of litterfall in secondary evergreen broadleaved forest. *Journal of Mountain Science*, 24(2): 215–221.
- Yang Y S, Xie J S, Chen Y X, He Z M, 2001. Amount, composition and seasonal dynamics of litterfall in mixed forest of *Cunninghamia lanceolata* and *Tsoongiodendron odorum*. *Scientia Silvae Sinicae*, 37(Suppl. 1): 30–34.
- Zhang D L, Mao Z J, Zhu S Y, Zhou B, 2008a. Litter falls of 6 major forest stands in Maoershan Mountain of Heilongjiang Province. *Bulletin of Botanical Research*, 28(1): 104–108.
- Zhang X M, Luo K L, Sun X Z, Tan J A, Lu Y L, 2006. Mercury in the topsoil and dust of Beijing city. *Science of the Total Environment*, 368(2-3): 713–722.
- Zhang X P, Wang X P, Zhu B, Zong Z J, Peng C H, Fang J Y, 2008b. Litter fall production in relation to environmental factors in northeast China's forests. *Journal of Plant Ecology*, 32(5): 1031–1040.