Response of bacterial *pdo1*, *nah*, and *C12O* genes to aged soil PAH pollution in a coke factory area

Xue-Mei Han · Yu-Rong Liu · Yuan-Ming Zheng ·
Xiao-Xia Zhang · Ji-Zheng He

Received: 23 September 2013 / Accepted: 15 April 2014 / Published online: 30 April 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract Soil pollution caused by polycyclic aromatic hydrocarbons (PAHs) is threatening human health and environmental safety. Investigating the relative prevalence of different PAH-degrading genes in PAH-polluted soils and searching for potential bioindicators reflecting the impact of PAH pollution on microbial communities are useful for microbial monitoring, risk evaluation, and potential bioremediation of soils polluted by PAHs. In this study, three functional genes, *pdo1*, *nah*, and *C12O*, which might be involved in the degradation of PAHs from a coke factory, were investigated by real-time quantitative PCR (qPCR) and clone library approaches. The results showed that the *pdo1* and *C12O* genes were more abundant than the *nah* gene in the soils. There was a significantly positive relationship between the *nah* or *pdo1* gene abundances and PAH content, while there was no correlation between *C12O* gene abundance and PAH content. Analyses of clone libraries showed that all the *pdo1* sequences were grouped into *Mycobacterium*, while all the *nah* sequences were classified into three groups: *Pseudomonas*, *Comamonas*, and *Polaromonas*. These results indicated that the abundances

of *nah* and *pdo1* genes were positively influenced by levels of PAHs in soil and could be potential microbial indicators reflecting the impact of soil PAH pollution and that *Mycobacterium* were one of the most prevalent PAHs degraders in these PAH-polluted soils. Principal component analysis (PCA) and correlation analyses between microbial parameters and environmental factors revealed that total carbon (TC), total nitrogen (TN), and dissolved organic carbon (DOC) had positive effects on the abundances of all PAH-degrading genes. It suggests that increasing TC, TN, and DOC inputs could be a useful way to remediate PAH-polluted soils.

Keywords Polycyclic aromatic hydrocarbon · Bioindicator · qPCR · PAH-degrading gene · Persistent organic pollutants · Soil pollution

Introduction

Polycyclic aromatic hydrocarbons (PAHs) are recalcitrant toxic pollutants (Couling et al. 2010), of which 16 are listed as priority pollutants by the US Environmental Protection Agency (USEPA) (Mackay et al. 1992). PAHs are distributed ubiquitously throughout the environment and are generated continuously from incomplete combustion and thermal decomposition processes of organic matter, especially of coal and oil (Johnsen et al. 2005). Human and ecosystem health are threatened by PAHs because of their toxic, mutagenic, and carcinogenic properties and high enrichment ratio within living organisms (Johnsen et al. 2005; Peng et al. 2008). PAHs are composed of two or more fused aromatic rings in various structural configurations, characterized by low aqueous solubility, high solid-water distribution ratios, and chemical stability. The aqueous solubility and, as a consequence, the bioavailability of PAHs decrease with increasing number of aromatic rings (Peng et al. 2008). In addition, many studies

Responsible editor: Robert Duran

X.-M. Han · Y.-R. Liu · Y.-M. Zheng · J.-Z. He (✉)
State Key Laboratory of Urban and Regional Ecology, Research
Center for Eco-Environmental Sciences, Chinese Academy of
Sciences, Beijing 100085, China
e-mail: jzhe@rcees.ac.cn

X.-M. Han
School of Resources and Environment, University of Jinan,
Jinan 250100, China

X.-X. Zhang
Key Laboratory of Microbial Resources Collection and Preservation,
Ministry of Agriculture, Beijing 100081, China

J.-Z. He
Melbourne School of Land and Environment, University of
Melbourne, Parkville 3010, Victoria, Australia

have demonstrated that the extractability and the bioavailability of PAHs decrease significantly with time in the aging processes (Northcott and Jones 2001; Luo et al. 2012). Most PAHs released into the environment will finally enter into the soil ecosystem, where they will impair human and other organisms through the food chain (Fismes et al. 2002). Therefore, it is of great importance to detect the effect of PAH pollution on certain bacterial populations associated with PAH degradation and seek appropriate bioindicators to reflect the impact of PAH pollution in aged industrial soils.

Soil microorganisms play a key role in energy flow, nutrient cycling, and organic matter turnover including biodegradation of organic pollutants. Although PAHs may undergo chemical oxidation and photolysis, in most cases, microbial degradation is still the major process involved in the dissipation of PAHs in the soil (Park et al. 1990). On the other hand, PAHs in the soil exert pressure on various microbes which inhabit it. Microorganisms are sensitive to even low levels of pollutants in soil and rapidly respond to soil pollution, so microbial parameters are usually considered as potential bioindicators of soil pollution (Sun et al. 2012).

Many microorganisms with the capability of degrading different PAHs have been isolated based on culture-dependent techniques (Bastiaens et al. 2000; Juhasz et al. 2000; Chang et al. 2011). However, it is well known that culture-independent approaches allow for a broader recognition of microbial populations responsible for PAH degradation in environments (Baldwin et al. 2003; Jurelevicius et al. 2012). Recently, the abundance and composition of PAH-degrading microbial populations present in PAH-polluted environments have been investigated to some extent through direct extraction of DNA from the environment and analysis of PAH-degrading genes (Singleton et al. 2005; Lillis et al. 2010; Paissé et al. 2012). *pdo1*, *nah*, and *C12O* are three important PAH-degrading genes which have been detected widely from environments polluted by aromatic compounds or bacteria isolated from those environments (Baldwin et al. 2003; Johnsen et al. 2006; Khan et al. 2009; Tuan et al. 2011). The *pdo1* gene encodes the pyrene dioxygenases associated with degradation of high-molecular weight (HMW) PAHs (Johnsen et al. 2006), the *nah* gene encodes naphthalene dioxygenases associated with degradation of low-molecular weight (LMW) PAHs (Baldwin et al. 2003), and the *C12O* gene encodes catechol 1,2-dioxygenase associated with cleavage of the last aromatic ring in the degradative pathway of PAHs (Sei et al. 1999). Johnsen et al. (2006) analyzed *nah*, *phnA*, and *pdo1* genes using a PCR assay but only detected the presence of *nah* and *pdo1* in soil microcosms spiked with street dust as a PAH source. Among the two genes detected, the *pdo1* gene was found to play an important role in the degradation of PAHs in the environment. Sei et al. (2004) found that abundances of the *C12O* and *C23O* genes increased significantly in aromatic compound spiked seawater

compared to those in the control, whereas community structure of bacterial 16S rDNA tended to be simplified by the spiked aromatic compound. To our knowledge, most of these studies focused on description of PAH-degrading genes based on microcosms spiked with a few PAHs (e.g., Scelza et al. 2007; Li et al. 2012; Meynet et al. 2012), while studies on comparison of genes encoding different PAH-degrading enzymes in field sites polluted by aged PAHs are scant. Therefore, this study aimed to investigate the prevalence of different PAH-degrading genes and the relationship between the PAH-degrading genes and levels of PAHs in soils from a coke factory area. The abundances of the *pdo1*, *nah*, and *C12O* genes in each soil were determined using real-time quantitative PCR (qPCR) assay, and the community structures of the PAH-degrading genes were analyzed by clone libraries and sequencing.

Materials and methods

Sampling site and soil collection

Soil samples classified as sandy loam were collected from a coke factory area. The coke factory was located in Xintai (35° 54' N, 117° 43' E), Shandong province of northern China. It started its operation in 2007 and can produce six million tons of secondary petroleum coke every year. Forty sampling sites were selected randomly in an area of about 16 ha around the coking oven. In each site, five soil subsamples were taken from the top 20 cm of the profile and mixed into one composite sample. All samples were placed on ice and transported to the laboratory, homogenized thoroughly after being gently crumbled to pass through a 2-mm sieve, and stored at −20 °C prior to use. Soil samples taken from the fridge were used directly for measurement of total water contents (TWC) and dissolved organic carbon (DOC) as well as extraction of total DNA. Air-dried soil samples passing through a 100-mesh sieve were used to determine the contents of total carbon (TC) and total nitrogen (TN). Freeze-dried soil samples passing through a 100-mesh sieve were used for extraction of PAHs.

Analysis of soil properties and PAH contents

TWC of the soil samples was determined through weight loss at 105 °C for 24 h. Soil pH was measured in a ratio of soil to water of 1:2.5 using a pH meter. TC and TN of the soil samples were determined using the Dumas method by an element analyzer (Vario EL III, Elementar, Hanau, Germany). DOC of the soil samples was measured following the method described by Sparling et al. (1998).

PAHs in the soil samples were extracted using ultrasonic extraction method (USEPA 2000). Briefly, the PAHs in 1 g of freeze-dried soil, which had been grounded and passed through a 100-mesh sieve, were extracted by 30 ml of

dichloromethane-hexane (1:1, v:v) in an ultrasonic bath for three times. The extract was collected by centrifugation and concentrated with a rotary evaporator. The concentrated extract was cleaned up with a chromatography column filled from bottom to top with 80-mesh neutral silica gel (12 cm, 180 °C heated for 12 h), 100-mesh neutral alumina (6 cm, 250 °C heated for 12 h), and anhydrous sodium sulfate (1 cm, 450 °C heated for 6 h). PAHs were eluted using 70 ml of hexane:dichloromethane (7:3, v:v) after alkanes removed by 15 ml of hexane. The concentrated eluent contain PAH mixture with a rotary evaporator continued to be concentrated to 1 ml through evaporation under a stream of N₂ gas.

PAHs were separated on an Agilent HP7890 gas chromatograph equipped with Agilent DB-5MS column (30 m×0.25 mm×0.25 µm) under a flow of 1 mL min⁻¹. The temperature program was as follows: 2 min at 80 °C, to 120 °C at 3 °C min⁻¹, to 200 °C at 5 °C min⁻¹, to 290 °C at 7 °C min⁻¹, and at 290 °C for 15 min. PAH detection was performed with an Agilent 5975C mass spectrometer with selected ion mode using electron impact at 70 eV.

Hexamethylbenzene was used as the internal standard and a PAH solution mix consisting of 16 USEPA priority PAHs (AccuStandard, New Haven, CT, USA) as the external standard. PAHs were identified on the basis of their retention times, target and qualifier ions, and quantified by a calibration curve (the ratio of external to internal standard). Phe-d10 was spiked into the soil samples 24 h prior to extracting PAHs to determine the recovery rate of the extraction method used.

Soil DNA extraction and qPCR assays

Total community DNA was extracted from 0.5 g soil using the MoBio Ultraclean soil DNA isolation kit (MoBio Laboratories, Solana Beach, CA, USA) according to the manufacturer's instructions. The concentration of the extracted DNA was determined using NanoDrop ND-1000 UV–vis spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA).

PAH-degrading genes, *nah*, *pdo1*, and *C12O*, were amplified on an iCycler iQ 5 thermocycler (Bio-Rad Laboratories, Hercules, CA, USA). The reaction mixture (25 µL) contained 2 µL of template DNA (1–10 ng), 0.5 µL of each of the forward and reverse primers (10 µM), and 12.5 µL of 2× SYBR[®] Premix Ex Taq[™] (TaKaRa Biotechnology, Dalian, China). The primer pairs and amplification conditions are shown in Table 1.

Standard curves for qPCR assays were constructed following the procedure described by Lillis et al. (2010). Briefly, *nah*, *pdo1*, and *C12O* genes were amplified from extracted DNA with the primer pairs listed in Table 1. The PCR products were cloned using the pGEM-T Easy Vector system (Promega Madison, WI, USA). Plasmids from the positive clones with each target gene insert were extracted and sequenced. The copy numbers of the plasmid DNA were calculated after determining the concentration. Serial dilutions were used to generate an external standard curve.

Construction of clone libraries and phylogenetic analysis

Clone libraries of the *pdo1* and *nah* genes were constructed because they were positively correlated with the total PAH content. The PCR products of the *pdo1* and *nah* genes from eight representative soil samples with different levels of PAHs were pooled and purified using a gel purification kit (Promega, Madison, WI, USA). The clone libraries of the genes were constructed by cloning PCR products using the pGEM-T Easy Vector system (Promega, Madison, WI, USA) according to the manufacturer's instructions, respectively. Eighty white colonies were randomly selected from each clone library, respectively. Inserts of each clone were amplified by PCR using the vector-specific primers T7 (5'-TAATACGACTCACTATAGGG-3') and SP6 (5'-TATTTAGGTGACACTATAG-3') following the method described by Lillis et al. (2010). Then, 50 and 74 positive clones from *pdo1* and *nah* clone libraries were used for sequencing, respectively.

Table 1 Primer pairs and amplification conditions used in real-time quantitative PCR (qPCR) assays

Target genes	Amplicon size (bp)	Primer pairs	Sequence (5'–3')	Amplification conditions	Reference
<i>nah</i>	377	NAH-F NAH-R	CAAAARCACCTGATTYATGG AYRCGRGSGACTTCTTTCAA	94 °C for 1 min (1 cycle); 94 °C for 20 s, 60 °C (reduced by 1 °C per cycle) for 30 s, 72 °C for 30 s (5 cycles); 94 °C for 20 s, 55 °C for 30 s, 72 °C for 30 s, and plate read at 83 °C for 15 s (35 cycles)	Baldwin et al. 2003
<i>pdo1</i>	793	pdo1-f pdo1-r	GTTCTACCTCGACCTCATTGCG CTGACCCATGTATTCCAGCC	94 °C for 5 min (1 cycle); 94 °C for 1 min, 60 °C (reduced by 0.5 °C per cycle) for 1 min, 72 °C for 1 min (10 cycles); 94 °C for 1 min, 50 °C for 1 min, 72 °C for 1 min, and plate read at 83 °C for 10 s (30 cycles)	Johnsen et al. 2006
<i>C12O</i>	324	C12O_F C12O_R	CGCGGATTGTNGAYST NTGGCANGCNAAYAC GACTCAGGTNGCRWANGC RAARTCRTC	95 °C for 1 min (1 cycle); 94 °C for 20 s, 60 °C (reduced by 1 °C per cycle) for 30 s, 72 °C for 30 s (5 cycles); 94 °C for 20 s, 55 °C for 30 s, 72 °C for 30 s (plate read simultaneously) (35 cycles)	Tuan et al. 2011

Classifications of the *pdo1* and *nah* sequences were identified by aligning with the BLAST search program (<http://blast.ncbi.nlm.nih.gov>). For each clone library, sequences sharing a 97 % nucleotide sequence identity were defined as one operational taxonomic unit (OTU) using MOTHUR software. Coverage of each clone library was computed according to the following equation: $C=1-(n/N)$, where n was the number of OTU containing only one clone and N was the total number of clones examined (Chelius and Triplett 2001). Only one representative sequence of each OTU was used for phylogenetic analyses. Phylogenetic trees were constructed using the neighbor-joining method with 1,000 bootstrap replicates of MEGA version 4.0. Sequences of the *pdo1* and *nah* clones used in phylogenetic tree construction were deposited in the GenBank nucleotide sequence database under accession numbers KF561983 to KF561985 and KC878841 to KC878848, respectively.

Statistical analyses

One-way analysis of variance (ANOVA) based on Fisher LSD test using SPSS 13.0 was used to assess differences between the abundances of different genes in the same soil. Principal component analysis (PCA) was conducted using CANOCO 4.5 to indicate the relationship between abundances of PAH-degrading genes and environmental factors including soil properties and content of PAHs. Spearman's correlation between various parameters was calculated using the STATISTIC 6.0. The probability levels of $P<0.05$ and $P<0.01$ were considered to be statistically significant.

Results

Soil basic properties and content of PAHs

Soil basic properties are shown in Table 2. Sample nos. were designated as nos. 1 to 40, respectively, from low to high content of PAHs in soils. The soil pH ranged from 4.44 to 7.97. The contents of TN, TC, and DOC ranged from 0.46 to 2.63 g kg⁻¹, from 6.15 to 120 g kg⁻¹, and from 19.3 to 43.8 mg kg⁻¹ of dry weight soil, respectively. The 16 USEPA priority PAHs were determined in this study. The recoveries for standard Phe-d10 spiked into the soil samples ranged from 70.0 to 94.5 %. The contents of total PAHs varied widely from 1.15 to 761 mg kg⁻¹ of dry weight soil (Table 2). The PAH mixture was divided into five groups according to their number of aromatic rings. Three- to four-ring PAHs were most common in all the soils, making up from 55.5 to 84.1 % of total PAHs (Fig. 1). The proportion of HMW PAHs increased with increasing content of total PAHs in soils, while LMW PAHs showed the opposite trend. Phenanthrene, fluoranthene, and pyrene were major compounds in all soils. Naphthalene was also one of the major components of

Table 2 Basic properties of the soil samples collected from a coke factory

Sample no. ^a	TWC (%)	pH	TN (g kg ⁻¹)	TC (g kg ⁻¹)	DOC (mg kg ⁻¹)	Total PAHs (mg kg ⁻¹)
1	11.6	7.88	0.46	6.15	22.7	1.15
2	9.70	7.70	0.55	6.28	22.5	1.53
3	9.40	7.56	0.54	6.29	22.6	1.54
4	15.7	7.54	1.00	13.2	25.4	2.18
5	7.70	7.92	0.47	6.69	24.5	2.24
6	9.70	7.65	0.56	7.08	22.9	2.32
7	15.3	7.61	0.82	9.43	24.9	2.62
8	16.7	7.39	0.80	10.7	27.4	2.75
9	14.9	6.81	0.87	15.3	26.7	3.02
10	15.1	5.87	1.06	10.5	23.1	3.59
11	15.3	7.55	1.05	15.2	24.8	3.61
12	16.2	7.73	0.86	11.8	24.6	3.71
13	15.0	5.87	0.84	9.22	20.5	4.38
14	15.7	6.28	0.74	9.54	26.1	4.53
15	14.3	7.69	0.93	13.3	23.3	5.93
16	20.2	4.44	1.41	52.9	28.5	6.35
17	13.3	6.39	0.97	14.5	31.1	6.92
18	18.5	6.73	1.48	52.7	34.2	6.99
19	10.6	5.87	1.34	26.1	39.7	8.76
20	18.1	7.71	1.70	63.3	34.7	10.2
21	18.4	7.25	0.98	26.7	25.5	10.2
22	17.5	6.57	1.27	57.3	26.3	11.7
23	16.5	6.56	0.98	41.3	25.8	12.6
24	15.4	5.99	1.90	81.0	33.7	12.7
25	16.2	7.66	0.76	18.9	28.7	13.2
26	15.6	7.70	0.80	17.6	19.3	13.4
27	14.6	7.66	0.88	18.7	26.7	13.8
28	13.4	7.71	0.97	24.8	26.6	13.9
29	12.1	6.89	1.77	32.1	33.1	14.1
30	12.5	6.60	1.81	78.8	32.1	14.3
31	14.8	5.22	1.71	28.3	30.5	15.9
32	16.2	7.60	1.25	42.0	37.9	21.2
33	15.3	7.34	1.69	33.4	34.4	22.4
34	9.80	7.70	1.36	40.2	29.2	22.8
35	12.7	7.56	0.92	36.9	24.6	26.1
36	13.1	6.38	2.63	120	42.5	30.1
37	10.6	7.70	1.27	40.7	28.2	30.5
38	14.4	7.97	1.37	44.2	30.4	33.5
39	15.7	7.72	0.87	22.5	28.2	113
40	13.0	7.60	1.65	61.1	43.8	761

TWC total water content, TN total nitrogen, TC total carbon, DOC dissolved organic carbon

^a Sample nos. were designated as nos. 1 to 40, respectively, from low to high content of PAHs in soils

PAHs in the soils polluted slightly by PAHs, where, five- to six-ring PAHs were almost undetectable.

Fig. 1 Compositional features of PAHs in the soils from a coke factory

Abundances of the *pdo1*, *nah*, and *C12O* genes

Copy numbers of the genes involved in PAH degradation are shown in Fig. 2. Abundance of the *pdo1* gene in soil samples except sample no. 26 and *C12O* gene in soil samples except sample nos. 21, 26, 39, and 40 was significantly higher than that of the *nah* gene ($P < 0.05$). Abundances of the *pdo1* and *C12O* genes were in the ranges of 1.63×10^6 – 1.08×10^9 and 8.68×10^6 – 7.67×10^8 copies g^{-1} of dry soil, respectively. However, the *nah* gene had the lowest copy number from 5.10×10^4 to 1.24×10^8 copies g^{-1} of dry soil. Abundances of the *pdo1* and *C12O* genes were 2.33–3,421 times and 0.54–4,000 times higher than that of the *nah* gene, respectively.

Diversity of the *pdo1* and *nah* gene sequences in the clone library

The clone library of *pdo1* gene had a coverage of 98.0 %. Fifty clones obtained from the library were assigned into three OTUs. The phylogenetic tree of the *pdo1* gene showed that all of OTUs were closely related to *Mycobacterium* (Fig. 3a). The coverage of *nah* gene clone library was computed to be 97.3 %. Seventy-four clone sequences of the library were assigned into eight OTUs. The phylogenetic tree of the *nah* gene assigned 45.9 % of the sequences to *Pseudomonas* of Gammaproteobacteria, 17.6 % to *Polaromonas* of Betaproteobacteria, and 36.5 % to *Comamonas* of Betaproteobacteria (Fig. 3b).

Fig. 2 Abundance of different PAH-degrading genes in the soils from a coke factory. Error bars are the SD of $n=3$

Fig. 3 Phylogenetic trees of *pdo1* (a) and *nah* (b) gene sequences retrieved from the soils in a coke factory. Designation of the clones included the following information: clone number, nucleotide sequence accession number, and amount of clones with similarities above 97 % to the listed sequence in the clone library in the parentheses. Numbers on nodes represent bootstrap values. The scale bar represents 0.2 % estimated sequence divergence

Relationship of the PAH-degrading gene abundances and environmental factors

PCA analysis of the PAH-degrading genes and environmental factors including soil properties and content of PAHs is illustrated in Fig. 4. PC1 and PC2 accounted for 67.1 and 27.1 % of the total variation, respectively. Of the genes involved in PAH degradation, the *nah* gene abundance had the closest relationship with the content of total PAHs. However, abundances of the other PAH-degrading genes, *pdo1* and *C12O*, were influenced more significantly by TC, TN, and DOC than PAHs. Table 3 showed that abundances of the *nah* and *pdo1* genes significantly correlated with the content of total PAHs ($R=0.76$ and 0.57 , respectively, $P<0.01$). However, the *C12O* gene had no significant relationship with content of total PAHs. All of the PAH-degrading genes were correlated significantly with TC (0.54 , 0.48 , and 0.48 for the *pdo1*, *nah*, and *C12O*, respectively, $P<0.01$), TN (0.69 , 0.42 , and 0.44 for the *pdo1*, *nah*, and *C12O*, respectively, $P<0.01$), and DOC (0.64 , 0.56 , and 0.34 for the *pdo1*, *nah*, and *C12O*, respectively, $P<0.01$ for the *pdo1* and *nah*, $P<0.05$ for the *C12O*). TWC and pH had no significant correlation with abundances of the *pdo1*, *nah*, and *C12O* genes.

Discussion

In this study, the collected soils from a coke factory were polluted by mostly three- to four-ring PAHs, especially

phenanthrene, fluoranthene, and pyrene. This result was similar to the previous report by Lors et al. (2010). They analyzed

Fig. 4 The principal component analyses (PCA) of PAH-degrading genes and environmental factors of the soils from a coke factory. The values in the parentheses represent the percentage of total variation explained by each axis

Table 3 Correlation coefficients between PAH-degrading gene abundance and environmental factors of the soils from a coke factory

	TWC	pH	TN	TC	DOC	PAHs	<i>pdo1</i>	<i>nah</i>	<i>C12O</i>
TWC	1.00								
pH	−0.32*	1.00							
TN	0.19	−0.40*	1.00						
TC	0.19	−0.28	0.88**	1.00					
DOC	0.06	−0.20	0.79**	0.71**	1.00				
PAHs	−0.06	0.12	0.21	0.23	0.46**	1.00			
<i>pdo1</i>	0.03	−0.16	0.69**	0.54**	0.64**	0.57**	1.00		
<i>nah</i>	−0.01	0.07	0.42**	0.48**	0.56**	0.76**	0.72**	1.00	
<i>C12O</i>	0.20	−0.15	0.44**	0.48**	0.34*	−0.08	0.09	−0.11	1.00

Significant correlations were presented in the form of italics

* $P < 0.05$; ** $P < 0.01$, significance level

the compositional features of PAH mixture in a coal tar contaminated soil and found that three- and four-ring PAHs were primarily and secondly abundant. Moreover, phenanthrene was the most concentrated compound among the three-ring PAHs, and fluoranthene and pyrene were the two most concentrated compounds among the four-ring PAHs. It indicated that soils polluted by the coking industry could be characterized by high proportions of three- and four-ring PAHs. Interestingly, the portion of HMW PAH increased with increasing total PAH content, while LMW PAHs showed a contrary trend in the examined soils. Some studies and the results here have shown that microbial populations involved in PAH degradation may be enriched by PAH pollution (Laurie and Lloyd-Jones 2000; Niepceon et al. 2013), and it is also well known that HMW PAHs are more resistant to degradation by soil microbes than LMW PAHs (Johnsen et al. 2005; Lers et al. 2010). So a higher proportion of LMW PAHs may be eliminated in seriously polluted soils than in slightly polluted soils, and consequently, HMW PAHs tend to accumulate in seriously polluted soils (Johnsen et al. 2006).

To elucidate the relationship between the abundances of various PAH-degrading genes and PAH content, the genes of *pdo1*, *nah*, and *C12O* which might be responsible for PAH degradation were quantified by qPCR in this study. Abundance of the *C12O* gene in soils was high but had no significant correlation to total PAH content. Catechol 1,2-dioxygenase encoded by the *C12O* gene catalyzes the oxygenative ring cleavage of catechol, which is the intermediate metabolite of many aromatic compounds including PAHs (Da Cunha et al. 2006). So, abundance of the *C12O* gene might be influenced by the content of not only total PAH but also many other kinds of aromatic compounds in soil. However, there were very significant positive correlations between the abundances of both *nah* and *pdo1* genes and total PAH content. Naphthalene dioxygenase (NDO) encoded by the *nah* gene has a relaxed substrate specificity and catalyzes the dioxygenation of many related two- and three-ring PAHs such as naphthalene, phenanthrene, anthracene, acenaphthylene, fluorene, and so on to their respective *cis*-dihydrodiols (Resnick et al. 1996). The *pdo1* gene sequences detected

showed high levels of similarity with *pdo1* genes identified from *Mycobacterium* which is involved in degradation of two-, three-, and four-ring PAHs including naphthalene, phenanthrene, fluoranthene, pyrene, chrysene, and benz(a)anthracene as well as even five-ring PAHs e.g., benz(a)pyrene with the presence of pyrene as co-metabolite (Schneider et al. 1996; Johnsen et al. 2006; Zhang et al. 2006). Thus, it was speculated that more organisms containing the *nah* and *pdo1* genes thrive under the condition of high PAH levels in soils. Similar results have been reported in recent studies (Laurie and Lloyd-Jones 2000; Jin et al. 2012). Laurie and Lloyd-Jones (2000) found that copy numbers of *phnAc* and *nahAc* genes involved in PAH degradation in PAH- and petroleum-polluted soils were significantly greater than those in a pristine soil using competitive PCR. Jin et al. (2012) found that naphthalene dioxygenase (NDO) genes were enhanced by naphthalene addition in laboratory microcosms. Johnsen et al. (2006) also found increased prevalence of *nah* and *pdo1* genes in soil microcosms spiked with street dust as a PAH source. These results indicated that the abundances of *nah* and *pdo1* genes were positively influenced by levels of PAHs and could be potential bioindicators reflecting the impact of PAH pollution on microbial communities in soil.

In all soils examined except sample no. 26, abundance of the *pdo1* gene was significantly higher than that of the *nah* gene. Although the approach based on DNA analysis did not reflect the relative contribution of these genes to degradation of PAHs, it did reveal the higher prevalence of bacterial populations harboring the *pdo1* gene than the *nah* gene. In this study, the clone library of the *pdo1* gene showed that all *pdo1* sequences were grouped into *Mycobacterium*, which was unsurprising as the primer used was designed based on the *pdo1* gene from *Mycobacterium* sp. strain 6PY1 (Krivobok et al. 2003; Johnsen et al. 2006). Johnsen et al. (2007) also found that *Mycobacteria* were dominant members of phenanthrene, pyrene, and fluoranthene degraders in soils originating from an asphalt and tar production plant. Thus, it was suggested that *Mycobacteria* may be one of the most prevalent PAH degraders in the soils examined. The dominance of *Mycobacteria* corresponded to the compositional

features of PAHs in the examined soils with a high share of three- to four-ring PAHs, especially phenanthrene, fluoranthene, and pyrene, which can be utilized and degraded by *Mycobacteria* harboring the *pdo1* gene (Johnsen et al. 2006). Some reports showed that *Mycobacteria* had a competitive advantage over PAH degraders from other bacterial genera, due to physiological adaptation to oligotrophy and low PAH bioavailability (Wick et al. 2002; Wells et al. 2005; Uyttenbroek et al. 2006).

The phylogenetic analysis showed that all sequences obtained from the *nah* gene clone library were classified into one of three groups: *Pseudomonas* of Gammaproteobacteria, *Comamonas* of Betaproteobacteria, and *Polaromonas* of Betaproteobacteria. Moreover, there was a very significant correlation between the *nah* gene abundance and PAH content. The close relationship between Gammaproteobacteria and Betaproteobacteria and PAH in soils has been reported in recent studies, especially Gammaproteobacteria has been suggested as a bioindicator of PAH pollution (Lors et al. 2010; Niepceon et al. 2013). Lors et al. (2010) monitored the PAH-degrading bacterial community structures by PCR and clone library in a coal-tar-contaminated soil in the process of wind-row treatment. Gammaproteobacteria were detected throughout the process, while Betaproteobacteria were detected when the accumulated content of two-, three-, and four-ring PAHs decreased substantially. Thus, the Gammaproteobacteria were suggested as a bioindicator of the potential biodegradation of PAHs in contaminated soil. Niepceon et al. (2013) monitored the abundances of ten bacterial phyla and classes by qPCR and *Pseudomonas* isolates using direct culture in soil microcosms spiked with a mixture of seven PAHs over a 90-day period and found that Betaproteobacteria and Gammaproteobacteria were more sensitive to the PAH mixture than other bacterial groups. Abundance of the Betaproteobacteria was significantly higher compared to the controls only at 60 days of incubation, while abundances of the Gammaproteobacteria and culturable *Pseudomonas* (affiliated to the Gammaproteobacteria) significantly increased over the longer period of incubation in response to the addition of PAH mixture. Thus, the Gammaproteobacteria was suggested by Niepceon et al. as a bioindicator to detect the impact of multiple pollution by PAHs on bacterial communities. In this study, the *nah* gene clone library displayed that the abundance levels of Gammaproteobacteria and Betaproteobacteria were 45.9 and 54.1 %, respectively. Although whichever of the two microbial groups is more sensitive cannot be determined according to that, it suggests that Gammaproteobacteria and Betaproteobacteria are the sensitive microbial taxa of soil PAH pollution.

PAH-degrading genes could be also influenced by soil characteristics. In this study, the content of TC, TN, and DOC were correlated significantly with abundance of the PAH-degrading genes e.g., *pdo1*, *nah*, and *C12O*. TC and

TN contents reflect organic nutrition in soils, and DOC represents the part of organic nutrition that could be utilized by microorganisms directly and easily (Cook and Allan 1992). So, soil microorganisms including PAH-degrading bacteria could thrive along with the increase of TC, TN, and DOC in soils (Yang et al. 2011; Zhang et al. 2012). Besides providing the nutrients for microorganisms, DOC may increase the solubility and bioavailability of PAHs (Thiele-Bruhn and Brümmer 2004; Kobayashi et al. 2008). For soils polluted by aged PAHs, the low bioavailability of PAHs was usually the key factor that restricted the degradation of PAHs by microorganisms (Kobayashi et al. 2008; Couling et al. 2010; Larsson et al. 2013). Therefore, it has been suggested that an effective method for remediation of PAH-polluted soils would be to apply organic substances thus increasing inputs of TC, TN, and DOC (Li et al. 2012).

Conclusions

The relationship between abundances of different PAH-degrading genes and content of PAHs in the soils from a coke factory was examined in this study. Abundances of the *nah* and *pdo1* genes were significantly correlated with the soil total PAH content. However, no significant relationship was found between the *C12O* gene abundance and the soil total PAH content. This indicated that abundances of the *nah* and *pdo1* genes may be potential microbial indicators reflecting the impact of PAH pollution on microbial communities in soil. Abundance of the *pdo1* gene was significantly higher than that of the *nah* gene in great majority of the soil samples. Analyses of clone libraries showed that all *pdo1* sequences were grouped into *Mycobacterium*, while all *nah* sequences could be classified into one of three groups: *Pseudomonas*, *Comamonas*, and *Polaromonas*. It suggested that *Mycobacteria* were more prevalent than bacterial taxa possessing the *nah* gene in aged PAH polluted soils. The significant positive correlations between contents of TC, TN, and DOC and abundances of all PAH-degrading genes implied that increasing inputs of TC, TN, and DOC into soils may be a useful way to remediate the soils polluted by PAHs. These findings provide new insights into microbial monitoring, risk evaluation, and potential bioremediation of soils polluted by the coking industry. However, more studies are still needed to confirm these conclusions in other soil types or environments polluted by aromatic compounds in different physicochemical conditions.

Acknowledgments This study was financially supported by the Natural Science Foundation of China (41201523, 41301567), the Natural Science Foundation of Shandong Province (ZR2011DL007), and the National High Technology Research and Development Program (863) of China (2012AA101403).

References

- Baldwin BR, Nakatsu CH, Nies L (2003) Detection and enumeration of aromatic oxygenase genes by multiplex and real-time PCR. *Appl Environ Microbiol* 69:3350–3358
- Bastiaens L, Springael D, Wattiau P, Harms H, DeWachter R, Verachtert H, Diels L (2000) Isolation of adherent polycyclic aromatic hydrocarbon (PAH)-degrading bacteria using PAH-sorbing carrier. *Appl Environ Microbiol* 66:1834–1843
- Chang CH, Lee J, Ko BG, Kim SK, Chang JS (2011) *Staphylococcus* sp. KW-07 contains *nahH* gene encoding catechol 2,3-dioxygenase for phenanthrene degradation and a test in soil microcosm. *Int Biodeterior Biodegrad* 65:198–203
- Chelius MK, Triplett EW (2001) The diversity of archaea and bacteria in association with the roots of *Zea mays* L. *Microbiol Ecol* 41:252–263
- Cook BD, Allan DL (1992) Dissolved organic carbon in old field soils: compositional changes during the biodegradation of soil organic matter. *Soil Biol Biochem* 24:595–600
- Couling NR, Towell MG, Semple KT (2010) Biodegradation of PAHs in soil: influence of chemical structure, concentration and multiple amendment. *Environ Pollut* 158:3411–3420
- Da Cunha CD, Rosado AS, Sebastian GV, Seldin L, von der Weid I (2006) Oil biodegradation by *Bacillus* strains isolated from the rock of an oil reservoir located in a deep-water production basin in Brazil. *Appl Microbiol Biotechnol* 73:949–959
- Fismes J, Perrin-Ganier C, Empereur-Bissonnet P, Morel JL (2002) Soil-to-root transfer and translocation of polycyclic aromatic hydrocarbons by vegetables grown on industrial contaminated soils. *J Environ Qual* 31:1649–1656
- Jin HM, Kim JM, Lee HJ, Madsen EL, Jeon CO (2012) *Alteromonas* as a key agent of polycyclic aromatic hydrocarbon biodegradation in crude oil-contaminated coastal sediment. *Environ Sci Technol* 46:7731–7740
- Johnsen AR, Wick LY, Harms H (2005) Principles of microbial PAH-degradation in soil. *Environ Pollut* 133:71–84
- Johnsen AR, de Liphay JR, Sørensen SJ, Ekelund F, Christensen P, Andersen O, Karlson U, Jacobsen CS (2006) Microbial degradation of street dust PAHs in microcosms simulating diffuse pollution of urban soil. *Environ Microbiol* 8:535–545
- Johnsen AR, Schmidt S, Hybholt TK, Henriksen S, Jacobsen CS, Andersen O (2007) Strong impact on the polycyclic aromatic hydrocarbon (PAH)-degrading community of a PAH-polluted soil but marginal effect on PAH degradation when priming with bioremediated soil dominated by *Mycobacteria*. *Appl Environ Microbiol* 73:1474–1480
- Juhasz AL, Stanley GA, Britz ML (2000) Microbial degradation and detoxification of high molecular weight polycyclic aromatic hydrocarbons by *Stenotrophomonas maltophilia* strain VUN 10,003. *Lett Appl Microbiol* 30:396–401
- Jurelevicius D, Alvarez WM, Peixoto R, Rosado AS, Seldin L (2012) Bacterial polycyclic aromatic hydrocarbon ring-hydroxylating dioxygenases (PAH-RHD) encoding genes in different soils from King George Bay, Antarctic Peninsula. *Appl Soil Ecol* 55:1–9
- Khan S, Heshan A, Qing G, Shuang L, He J (2009) Biodegradation of pyrene and catabolic genes in contaminated soils cultivated with *Lolium multiflorum* L. *J Soils Sediments* 9:482–491
- Kobayashi T, Navarro RR, Tatsumi K, Iimura Y (2008) Influence of compost amendment on pyrene availability from artificially spiked soil to two subspecies of *Cucurbita pepo*. *Sci Total Environ* 404:1–9
- Krivobok S, Kuony S, Meyer C, Louwagie M, Willison JC, Jouanneau Y (2003) Identification of pyrene-induced proteins in *Mycobacterium* sp. strain 6PY1: evidence for two ring-hydroxylating dioxygenases. *J Bacteriol* 185:3828–3841
- Larsson M, Hagberg J, Rotander A, van Bavel B, Engwall M (2013) Chemical and bioanalytical characterisation of PAHs in risk assessment of remediated PAH-contaminated soils. *Environ Sci Pollut Res*. doi:10.1007/s11356-013-1787-6
- Laurie AD, Lloyd-Jones G (2000) Quantification of *phnAc* and *nahAc* in contaminated New Zealand soils by competitive PCR. *Appl Environ Microbiol* 66:1814–1817
- Li XZ, Wu YC, Lin XG, Zhang J, Zeng J (2012) Dissipation of polycyclic aromatic hydrocarbons (PAHs) in soil microcosms amended with mushroom cultivation substrate. *Soil Biol Biochem* 47:191–197
- Lillis L, Clipson N, Doyle E (2010) Quantification of catechol dioxygenase gene expression in soil during degradation of 2,4-dichlorophenol. *FEMS Microbiol Ecol* 73:363–369
- Lors C, Ryngaert A, Périé F, Diels L, Damidot D (2010) Evolution of bacterial community during bioremediation of PAHs in a coal tar contaminated soil. *Chemosphere* 81:1263–1271
- Luo L, Lin S, Huang HL, Zhang SZ (2012) Relationships between aging of PAHs and soil properties. *Environ Pollut* 170:177–182
- Mackay D, Shiu WY, Ma KC (1992) Illustrated handbook of physical-chemical properties and environmental fate for organic chemicals, vol 2. Lewis Publishers, Michigan
- Meynet P, Hale SE, Davenport RJ, Cornelissen G, Breedveld GD, Werner D (2012) Effect of activated carbon amendment on bacterial community structure and functions in a PAH impacted urban soil. *Environ Sci Technol* 46:5057–5066
- Niepceron M, Martin-Laurent F, Crampon M, Portet-Koltalo F, Akpa-Vinceslas M, Legras M, Bru D, Bureau F, Bodilis J (2013) GammaProteobacteria as a potential bioindicator of a multiple contamination by polycyclic aromatic hydrocarbons (PAHs) in agricultural soils. *Environ Pollut* 180:199–205
- Northcott GL, Jones KC (2001) Partitioning, extractability, and formation of nonextractable PAH residues in soil. 1. Compound differences in aging and sequestration. *Environ Sci Technol* 35:1103–1110
- Paissé S, Goñi-Urriza M, Stalder T, Budzinski H, Duran R (2012) Ring-hydroxylating dioxygenase (RHD) expression in a microbial community during the early response to oil pollution. *FEMS Microbiol Ecol* 80:77–86
- Park KS, Sims RC, Dupont RR, Doucette WJ, Matthews JE (1990) Fate of PAH compounds on two soil types: influence of volatilization, abiotic loss and biological activity. *Environ Toxicol Chem* 9:187–195
- Peng RH, Xiong AS, Xue Y, Fu XY, Gao F, Zhao W, Tian YS, Yao QH (2008) Microbial biodegradation of polyaromatic hydrocarbons. *FEMS Microbiol Rev* 32:927–955
- Resnick SM, Lee K, Gibson DT (1996) Diverse reactions catalyzed by naphthalene dioxygenase from *Pseudomonas* sp strain NCIB 9816. *J Ind Microbiol* 17:438–457
- Scelza R, Rao MA, Gianfreda L (2007) Effects of compost and of bacterial cells on the decontamination and the chemical and biological properties of an agricultural soil artificially contaminated with phenanthrene. *Soil Biol Biochem* 39:1303–1317
- Schneider J, Grosser R, Jayasimhulu K, Xue W, Warshawsky D (1996) Degradation of pyrene, benz[a]anthracene, and benzo[a]pyrene by *Mycobacterium* sp. strain RJGII-135, isolated from a former coal gasification site. *Appl Environ Microbiol* 62:13–19
- Sei K, Asano K, Tateishi N, Mori K, Ike M, Fujita M (1999) Design of PCR primers and gene probes for the general detection of bacterial populations capable of degrading aromatic compounds via catechol cleavage pathways. *J Biosci Bioeng* 88:542–550
- Sei K, Inoue D, Wada K, Mori K, Ike M, Kohno T, Fujita M (2004) Monitoring behaviour of catabolic genes and change of microbial community structures in seawater microcosms during aromatic compound degradation. *Water Res* 38:4405–4414
- Singleton DR, Powell SN, Sangaiah R, Gold A, Ball LM, Aitken MD (2005) Stable-isotope probing of bacteria capable of degrading salicylate, naphthalene, or phenanthrene in a bioreactor treating contaminated soil. *Appl Environ Microbiol* 71:1202–1209
- Sparling GP, Vojvodić-Vuković M, Schipper LA (1998) Hot-water-soluble C as a simple measure of labile soil organic matter: the

- relationship with microbial biomass C. *Soil Biol Biochem* 30:1469–1472
- Sun FL, Wang YS, Sun CC, Peng YL, Deng C (2012) Effects of three different PAHs on nitrogen-fixing bacterial diversity in mangrove sediment. *Ecotoxicology* 21:1651–1660
- Thiele-Bruhn S, Brümmer GW (2004) Fractionated extraction of polycyclic aromatic hydrocarbons (PAHs) from polluted soils: estimation of the PAH fraction degradable through bioremediation. *Eur J Soil Sci* 9:567–578
- Tuan NN, Hsieh HC, Lin YW, Huang SL (2011) Analysis of bacterial degradation pathways for long-chain alkylphenols involving phenol hydroxylase, alkylphenol monooxygenase and catechol dioxygenase genes. *Bioresour Technol* 102:4232–4240
- USEPA (2000) Method 3550C, test methods for evaluating solid waste. EPA SW-846, 3rd edn. US Government Print Office, Washington
- Uytendaele M, Breugelmans P, Janssen M, Wattiau P, Joffe B, Karlson U, Ortega-Calvo JJ, Bastiaens L, Ryngaert A, Hausner M, Springael D (2006) Distribution of the *Mycobacterium* community and polycyclic aromatic hydrocarbons (PAHs) among different size fractions of a long-term PAH-contaminated soil. *Environ Microbiol* 8:836–847
- Wells M, Wick LY, Harms H (2005) Model polymer release system study of PAH bioaccessibility: the relationship between rapid release and bioaccessibility. *Environ Sci Technol* 39:1055–1063
- Wick LY, Ruiz de Munain A, Springael D, Harms H (2002) Responses of *Mycobacterium* sp. LB501T to the low bioavailability of solid anthracene. *Appl Microbiol Biotechnol* 58:378–385
- Yang Y, Zhang N, Xue M, Lu ST, Tao S (2011) Effects of soil organic matter on the development of the microbial polycyclic aromatic hydrocarbons (PAHs) degradation potentials. *Environ Pollut* 159:591–595
- Zhang XX, Cheng SP, Zhu CJ, Sun SL (2006) Microbial PAH-degradation in soil: degradation pathways and contributing factors. *Pedosphere* 16:555–565
- Zhang J, Lin XG, Liu WW, Wang YM, Zeng J, Chen H (2012) Effect of organic wastes on the plant-microbe remediation for removal of aged PAHs in soils. *J Environ Sci-China* 24:1476–1482