Polychlorinated biphenyls in sediments and fish species from the Murchison Bay of Lake Victoria, Uganda

Patrick Ssebugere^{a,c}, Mika Sillanpää^b, Pu Wang^c, Yingming Li^c, Bernard T. Kiremire^a, Gabriel N. Kasozi^a, Chaofei Zhu^c, Daiwei Ren^c, Nali Zhu^c, Haidong Zhang^c, Hongtao Shang^c, Qinghua Zhang^{c,*}, Guibin Jiang^c

^a Department of Chemistry, Makerere University, P.O. Box 7062, Kampala, Uganda

^b Laboratory of Green Chemistry, Lappeenranta University of Technology, Sammonkatu 12, 50130 Mikkeli, Finland

^c State Key Laboratory of Environmental Chemistry and Ecotoxicology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

HIGHLIGHTS

- High concentrations of PCBs were found at sites near wastewater discharges.
- The PCB concentrations were low to moderate compared to other locations worldwide.
- Based on the European Commission set TEQ, the fish were fit for human consumption.

ARTICLE INFO

Article history:

Received 12 December 2013

Received in revised form 2 March 2014

Accepted 2 March 2014

Available online 21 March 2014

Keywords:

Polychlorinated biphenyls

Sediments

Fish

Murchison Bay

Lake Victoria

ABSTRACT

Polychlorinated biphenyls (PCBs) were determined in sediments and two fish species collected from the Murchison Bay in Lake Victoria, using high resolution gas chromatography coupled to a high resolution mass spectrometer. Total PCB concentrations (Σ_{18} PCBs) varied widely with mean values ranging from 777 to 4325 pg g^{-1} dry weight (dw) for sediments and 80 to 779 pg g^{-1} wet weight (ww) for fish. The PCB levels in the sediments were significantly higher at the station closest to Nakivubo channel, presumably due to effluents discharged by the channel, which may contain domestically produced commercial PCB mixtures. For fish, the concentrations in Nile perch (*Lates niloticus*) were significantly greater than those in Nile tilapia (*Oreochromis niloticus*) at all study stations, possibly due to dietary differences among species. World Health Organization-toxic equivalents (WHO₂₀₀₅-TEQs) for the dioxin-like PCBs were 0.04–0.64 pg g^{-1} dw and 0.01–0.39 pg g^{-1} ww for sediments and fish, respectively. The non-ortho PCBs exhibited the highest contribution to the Σ_{12} TEQs (>75%) compared to the mono-ortho PCBs in both fish species. The TEQs in the present study were lower than many reported worldwide in literature for fish and were within the permissible level recommended by the European Commission, implying that the fish did not pose health hazards related to PCBs to the consumers.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Lake Victoria is a vital natural aquatic resource in Uganda, but its ecological health is being threatened, mainly by rapid urbanization and industrialization. These developments coupled with a population of over one million people in its catchment area have resulted in increased pollution (Banadda et al., 2009, 2011a, 2011b; Wasswa et al., 2010). Polychlorinated biphenyls (PCBs) have previously been documented in the lake (Ssebugere et al. 2013a, b; Ssebugere et al., 2014). The PCBs are released into the environment as a result of industrial applications such as capacitors, hydraulic fluids in electronic

equipment, and coolants and lubricants in transformers (Gidarakos et al., 2009; Moon et al., 2012; Lundin et al., 2013; Miller et al., 2013).

Exposure to PCBs has been associated with diseases like diabetes mellitus, cancer and hypertension (Lee et al., 2007a). Other adverse effects caused by the pollutants include endocrine disruption, neuro-behavioral deficits and fetal toxicity (Lee et al., 2007b; Ha et al., 2007; 2009; Uemura et al., 2009). The PCBs undergo long-range transport causing global pollution (Wiberg et al., 2009; Sundqvist et al., 2009a, b). Furthermore, the contaminants can accumulate in sediments close to point sources and in benthic organisms and fish due to their hydrophobic nature, low metabolic transformation and long half-lives (Suedel et al., 1994; Jönsson et al., 2003; Åberg et al., 2008; 2010; Lavandier et al., 2013). Fish can be an important monitoring tool, because they concentrate PCBs directly from water through their diet, and indicate hazards to piscivorous predators and humans (Van

* Corresponding author. Tel.: +86 13693263173.

E-mail address: qhzhzhang@rcees.ac.cn (Q. Zhang).

der Oost et al., 2003; Huang et al., 2006; Gustavson et al., 2008; Parnell et al., 2008).

To our knowledge whereas PCBs have been reported in fish and sediments from the Napoleon Gulf and Thurston Bay of Lake Victoria (Ssebugere et al., 2013a, b; Ssebugere et al., 2014) and from other water bodies in the world (Manirakiza et al., 2002; Adu-Kumi et al., 2010; Darko et al., 2008; Helm et al., 2008; Eqani et al., 2013; Verhaert et al., 2013), literature shows no work about these pollutants in the Murchison Bay. The Bay is a major hotspot for discharge of industrial outflow, urban runoff and municipal effluents from Kampala, Uganda's capital city (Banadda et al., 2009). The objectives of this study were to investigate the occurrence and contamination levels of PCBs in sediments and fish from the Murchison Bay.

2. Materials and methods

2.1. Study area

The area of study was the Murchison Bay located in the northwestern part of Lake Victoria and southeast of Kampala city (Fig. 1). The Bay covers an area of 60 km² and includes semi-enclosed inner and outer parts. The Murchison Bay is the main recipient of sewage effluents, industrial- and municipal-waste from the central district of Kampala City, via the Nakivubo channel. The channel passes through dense residential settlements and commercial areas. In the past, the channel ended in wetland areas, allowing wastewater to be drained into papyrus swamps before entering the Bay. However, in recent years the wetlands which played the role of secondary treatment have been degraded due to settlement and commercial activities (Wasswa, 2009). The water turbidity in the Bay has increased while the water transparency has declined due to excessive waste input.

2.2. Sample collection

A total of 24 surface sediments (<25 cm depth) were taken using a sediment corer from four stations namely 1 and 2 in the inner Bay, and 3 and 4 in the outer Bay (Fig. 1) in March, 2013. Six sediments were randomly taken from each station at distances of approximately

200 m from one another. The geographical coordinates of the inner and outer parts of the Murchison Bay are 00°15'72"N, 32°38'74"E and 00°08'71"N, 32°37'58"E, respectively. Ninety six fish of different species namely: Nile Perch (*Lates niloticus*) a carnivorous top-predator and Nile Tilapia (*Oreochromis niloticus*) a detritivorous species were collected using gill nets within the same locality as the sediments. These two species exhibit different feeding habitats and thus may be exposed differently to contamination through trophic transfer. The weights of *O. niloticus* ranged from 289 to 652 g (mean 389 g) while those of *L. niloticus* varied from 1628 to 2979 g (mean 2378 g). The mean lengths ($\pm$ standard deviation) were 29 ± 3 cm, and 68 ± 5 cm, respectively. The muscle tissues were dissected between the pectoral fin and vent of the fish, minced into pieces and subsamples taken. Subsamples of 4 fish from the same location and, of similar length and species were pooled and homogenized. The sediments and homogenized fish were transferred into acetone rinsed glass bottles and shipped to the Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences where they were freeze-dried.

2.3. Analytical method

Sample extraction and cleanup procedures followed the method reported by Wang et al. (2013). Briefly, 2 g of the freeze-dried sample (fish or sediment) was ground with 10 g of anhydrous sodium sulfate to a free floating powder using a mortar and pestle. The mixture was spiked with ¹³C-labeled surrogate standards (US EPA defined 68A-LCS) and extracted on an accelerated solvent extractor device (ASE 300, Dionex, USA) using 1:1 v/v mixture of *n*-hexane/dichloromethane (DCM). The resultant extract was concentrated to 2 mL on a rotary evaporator. The concentrated extract was divided into two subsamples for fish (0.3 and 1.7 mL). The 0.3 mL was used for gravimetric determination of the lipid content while the 1.7 mL was kept for clean-up. For the sediments, activated copper granules were added to the extract to remove elemental sulfur before purification with column chromatography. The extracts were then cleaned up with a multilayered silica column (packed from bottom to top with 1 g silica gel, 4 g basic silica gel, 1 g silica gel, 8 g acid silica gel, 2 g silica gel and 2 g anhydrous sodium sulfate), an alumina column (6 g basic alumina and 3 g anhydrous sodium sulfate

Fig. 1. Map of the Murchison Bay showing the study stations. Adapted from Haande et al. (2011).

Table 1

Mean concentrations and ranges (pg g⁻¹ dry weight) of PCBs in sediments from the Murchison Bay.

Congener	Station 1	Station 2	Station 3	Station 4
CB 28	375 ± 25 353–402	318 ± 30 284–340	103 ± 54 51–158	63 ± 21 45–85
CB 52	507 ± 99 397–589	293 ± 31 259–319	169 ± 14 153–179	108 ± 44 58–137
CB 101	585 ± 234 420–853	343 ± 176 229–546	158 ± 68 105–235	122 ± 41 75–150
CB 138	933 ± 446 631–1445	553 ± 396 230–1009	240 ± 131 151–390	153 ± 87 79–249
CB 153	898 ± 240 721–1171	585 ± 465 251–1115	227 ± 90 173–331	177 ± 85 110–272
CB 180	218 ± 63 152–276	117 ± 54 61–169	49 ± 23 26–72	29 ± 6.1 23–35
Σ indicator-PCBs	3516 2749–4652	2209 1464–3488	946 719–1282	652 409–921
CB 77	82 ± 20 60–99	56 ± 6.3 52–64	12 ± 4.5 6.3–14	8.8 ± 1.2 8.1–10
CB 81	1.9 ± 1.1 0.7–2.8	2.5 ± 0.9 1.6–3.3	1.0 ± 0.6 0.6–1.4	0.8 ± 0.3 0.6–1.0
CB 126	4.9 ± 1.4 3.3–5.8	2.2 ± 0.7 1.7–3.0	1.2 n.d.–1.2	0.9 ± 0.5 n.d.–1.2
CB 169	n.d.	0.3 n.d.–0.3	n.d.	0.1 n.d.–0.1
Σ non-ortho-PCBs	89 66–105	61 55–70	14 6.8–17	11 ± 2.0 8.9–13
CB 105	140 ± 70 99–221	59 ± 18 47–79	26 ± 10 19–38	22 ± 13 11–36
CB 114	6.7 ± 3.3 4.4–11	2.9 ± 0.6 2.3–3.4	1.9 ± 0.6 1.5–2.6	1.2 ± 0.6 0.7–1.9
CB 118	434 ± 214 302–682	174 ± 74 139–228	110 ± 59 71–177	72 ± 45 32–120
CB 123	53 ± 19 42–76	24 ± 7.4 18–32	13 ± 5.3 8.7–19	7.4 ± 6.2 1.0–13
CB 156	54 ± 28 36–86	28 ± 14 16–43	14 ± 5.1 9.2–19	7.5 ± 3.3 4.9–11
CB 157	7.7 ± 5.2 4.2–14	3.4 ± 1.3 2.2–4.8	2.0 ± 1.0 1.3–3.1	1.3 ± 0.7 0.8–2.1
CB 167	16 ± 7.9 11–26	8.9 ± 5.0 4.6–14	4.4 ± 1.1 3.1–5.2	1.9 ± 0.4 1.5–2.2
CB 189	8.0 ± 1.9 6.6–10	5.5 ± 2.7 3.0–8.4	2.7 ± 1.3 1.3–3.9	1.1 ± 0.4 0.8–1.6
Σ mono-ortho-PCBs	720 508–1124	306 231–413	174 124–266	114 53–189
Σ ₁₂ dl-PCBs	809 613–1190	367 289–482	188 131–283	125 62–202
Σ ₁₈ PCBs	4325 3762–5842	2576 1753–3970	1134 850–1365	777 471–1123
WHO ₂₀₀₅ -TEQs	0.55 0.37–0.64	0.25 0.19–0.33	0.11 0.04–0.12	0.10 0.07–0.14

The number of sediment samples per station (N) = 6; n.d. – non-detectable.

at the top) and carbon column (1.5 g of 18% carbon dispersed in cilite, 3 g anhydrous sodium sulfate at the top), respectively. The clean eluate was concentrated to 1 mL using a rotary evaporator and transferred into vials containing 20 µL nonane. The contents in the vials were further dried to 25 µL using a gentle stream of nitrogen and later spiked with injection standards (US EPA defined 68A-IS) prior to analysis.

2.4. Instrumental analysis

The analysis of 18 PCB congeners including 6 indicator PCBs (IUPAC numbers 28, 52, 101, 138, 153 and 180) and twelve dioxin-like PCBs (IUPAC numbers 77, 81, 126, 169, 105, 114, 118, 123, 156, 157, 167 and 189) was done using high-resolution gas chromatography coupled with high-resolution mass spectrometry (HRGC/HRMS) (AutoSpec Ultima, Waters, USA). Chromatographic separation was achieved by injecting 1 µL of sample on a fused silica capillary column (DB5MS, 60 m × 0.25 mm i.d. × 0.25 µm film thicknesses). The oven temperature of the GC was held at 120 °C for 1 min, then increased to 150 °C at a rate of 30 °C min⁻¹ and finally to 300 °C at 2.5 °C min⁻¹. Helium was used as a carrier gas at a flow rate of 1.0 mL min⁻¹. The HRMS was operated in VSIR mode at a resolution of ≥ 10000 under positive electron ionization conditions (35 eV) and a source temperature of 270 °C.

2.5. Quality assurance and quality control

To ensure the quality of data, surrogate PCB standards, blanks, replicates and a certified reference material (CRM) were included in the analysis. Recoveries for majority of the surrogate standards met the requirements of US EPA methods 1668A (were in the range of 75–127%) except for CBs 28, 118, and 138 whose recoveries were <75% (range: 56–74%). The recoveries resulting from triplicate determinations (N = 3) of the certified reference material were between 82% and 106%. The limit of detection (LOD) which was calculated as three times the signal-to-noise ratio, varied from 0.04 to 0.96 pg g⁻¹ dry weight (dw) for sediments and 0.02 to 0.80 pg g⁻¹ wet weight (ww) for fish. PCBs in the field and procedural blanks were detected at <15% of the concentration in each batch of 12 samples, so the reported data were not blank corrected.

2.6. Statistical data analysis

Statistical analysis was performed using Statistica® 7 for Windows. Kolmogorov–Smirnov test was used for verifying the normality of data. Since concentration values were not normally distributed, non-parametric statistics was applied. The Mann–Whitney U-test was used for pair wise comparison and the Kruskal–Wallis test when more than two groups were considered. In these two non-parametric tests, statistical significance was set at $p < 0.05$. In the present study, toxic

Table 2

Comparison of sediment PCB concentrations (pg g⁻¹ dry weight) in this study with those from other locations.

Area	PCBs (pg g ⁻¹ dry weight)	References
Napoleon Gulf and Thurston Bay of Lake Victoria, Uganda	64–1262 (18 congeners)	Ssebugere et al. (2013b) and Ssebugere et al. (2014)
Nile River, Egypt	1461–2244 (18 congeners)	El-Kady et al. (2007)
Pangani River basin, Tanzania	357–11000 (28 congeners)	Hellar-Kihampa et al. (2013)
Congo River basin, Democratic Republic of Congo	n.d.–1400 (33 congeners)	Verhaert et al. (2013)
Ghar El Melh lagoon, Tunisia	n.d.–3987 (20 congeners)	Ameur et al. (2011)
Mekong River delta, Vietnam	110–2000 (13 congeners)	Carvalho et al. (2008)
Riverine and coastal waters of Surabaya, Indonesia	n.d.–420000 (62 congeners)	Ilyas et al. (2011)
Scheldt River, Belgium	n.d.–200000 (27 congeners)	Covaci et al. (2005)
Remote lakes and coastal areas, Norwegian Arctic	180–13000 (15 congeners)	Jiao et al. (2009)
Mersey Estuary, United Kingdom	36000–1409000 (7 congeners)	Vane et al. (2007)
Lake Maggiore, Italy and Switzerland	n.d.–3000 (12 congeners)	Vives et al. (2007)
Besòs River, Barcelona	6–14 (30 congeners)	Castells et al. (2008)
Indiana Harbor and Ship Canal, Lake Michigan, United States of America	53000–35000000 (163 congeners)	Martinez et al. (2010)
Sea Lots, Port of Spain, Trinidad and Tobago	62000–601000 (136 congeners)	Mohammed et al. (2011)

n.d. – non-detectable.

Fig. 2. Percent contribution of the congeners to the Σ_{18} PCBs per station.

Fig. 3. Mean concentrations of the PCB congeners in the sediments per station.

equivalents (TEQs) for the 12 dioxin-like (dl) PCBs were calculated using individual congener concentrations and the 2005 toxic equivalency factors (TEFs) published by the World Health Organization (WHO) (van den Berg et al., 2006).

3. Results and discussion

3.1. Concentrations of PCBs in sediments from the Murchison Bay

Total PCB concentration (Σ_{18} PCBs) in Murchison Bay sediments varied widely with mean values ranging from 777 to 4325 pg g⁻¹ dw (Table 1). They were 3-fold higher than those reported in sediments from the Napoleon Gulf and Thurston Bay of Lake Victoria (Ssebugere et al., 2013b; Ssebugere et al., 2014). The PCB concentrations in the present study were comparable to data from other fresh water bodies in Africa: the Nile River in Egypt (El-Kady et al., 2007), Ghar El Melh lagoon in Tunisia (Ameur et al., 2011), Pangani River basin in Tanzania

(Hellar-Kihampa et al., 2013) and Congo River basin in the Democratic Republic of Congo (Verhaert et al., 2013). The results were also in the range of data found in sediments from tropical areas in Asia: the Mekong River delta in Vietnam (Carvalho et al., 2008) and riverine and coastal waters of Surabaya in Indonesia (Ilyas et al., 2011). The levels were lower compared to concentrations reported for industrialized areas in Europe and United States of America (Covaci et al., 2005; Vane et al., 2007; Vives et al., 2007; Castells et al., 2008; Jiao et al., 2009; Martinez et al., 2010; Mohammed et al., 2011) (Table 2).

The proportion of the 6 indicator PCBs (Σ_6 PCBs) to Σ_{18} PCBs in this study were between 80 and 88% (with a mean value of 84%). CB 138 and 153 were the predominant congeners, collectively accounting for 21% of the Σ_{18} PCBs on average (Fig. 2). Barakat et al. (2012) also reported the predominance of CB 138 and CB 153 in surface sediments from Lake Maryut in Egypt. The accumulation of PCBs in the environment is directly linked to their degree of chlorination, stereochemistry and lipophilicity (Mazet et al., 2005). PCBs with a high degree of chlorination

Table 3
Mean concentrations and ranges (pg g⁻¹ wet weight) of PCBs in fish species from the Murchison Bay.

	Station 1		Station 2		Station 3		Station 4	
	<i>L. niloticus</i>	<i>O. niloticus</i>	<i>L. niloticus</i>	<i>O. niloticus</i>	<i>L. niloticus</i>	<i>O. niloticus</i>	<i>L. niloticus</i>	<i>O. niloticus</i>
Lipid content (%)	2.8	1.5	2.3	1.4	2.2	1.1	1.6	1.2
<i>Congener</i>								
CB 28	38 ± 22 13–53	14 ± 7.0 6.0–19	62 ± 21 40–83	13 ± 3.6 8.4–15	23 ± 7.5 18–31	32 ± 14 12–68	26 ± 15 10–41	9.3 ± 0.7 8.6–10
CB 52	39 ± 23 23–65	30 ± 24 7.8–55	60 ± 52 10–128	22 ± 18 9.0–43	19 ± 8.2 10–27	13 ± 4.6 10–18	12 ± 3.2 9.6–16	8.1 ± 2.1 5.6–9.5
CB 101	58 ± 11 45–66	23 ± 18 2.1–37	86 ± 39 50–128	17 ± 4.9 12–22	50 ± 25 22–65	13 ± 2.6 10–15	21 ± 6.5 16–28	9.8 4.9–14
CB 138	183 ± 98 72–259	43 ± 9.9 32–51	84 ± 38 52–127	17 ± 13 7.7–32	61 ± 31 25–81	10 ± 1.3 9.8–12	62 ± 32 33–96	18 ± 13 5.9–31
CB 153	220 ± 58 75–386	37 ± 13 22–49	98 ± 55 51–158	22 ± 14 13–39	76 ± 43 30–112	12 ± 2.1 10–14	72 ± 15 37–133	18 ± 12 4.9–30
CB 180	98 ± 57 37–151	12 ± 10 5.1–24	8.3 ± 2.2 n.d.–9.8	3.8 ± 2.5 1.1–7.7	13 ± 6.7 8–21	3.2 ± 1.4 1.7–4.5	11 ± 6.7 6.0–18	4.9 ± 3.0 1.5–7.3
Σ indicator-PCBs	636 363–904	159 75–228	398 288–573	95 57–124	242 144–307	83 60–122	204 112–322	68 32–100
CB 77	5.5 ± 1.7 3.8–7.1	3.1 ± 2.5 1.2–5.9	n.d. n.d.–1.4	1.0 ± 0.4 0.6–1.4	2.2 ± 0.9 1.3–3.1	1.3 ± 0.2 1.0–1.4	1.3 ± 0.1 n.d.–1.4	0.7 n.d.–0.7
CB 81	0.9 n.d.–0.9	1.1 ± 0.6 n.d.–1.5	n.d.	n.d.	0.9 ± 0.2 n.d.–1.5	n.d.	0.5 ± 0.4 n.d.–0.8	1.1 ± 0.1 n.d.–1.1
CB 126	1.0 ± 0.7 0.5–1.5	n.d.	0.2 n.d.–0.2	n.d.	n.d.	n.d.	0.8 n.d.–0.8	n.d.
CB 169	n.d.	0.2 n.d.–0.2	n.d.	n.d.	n.d.	n.d.	0.2 n.d.–0.2	0.1 n.d.–0.1
Σ non-ortho PCBs	7.4 5.7–7.6	4.4 2.4–6.6	0.2 n.d.–0.2	1.0 0.6–1.4	3.1 2.4–3.1	1.3 1.0–1.4	2.8 n.d.–3.2	1.9 0.7–1.1
CB 105	23 ± 11 13–34	7.8 ± 1.8 5.8–8.9	1.8 ± 0.4 1.4–2.2	1.9 ± 0.3 1.7–2.2	3.6 ± 0.1 3.5–3.7	1.8 ± 0.8 1.0–2.6	6.7 ± 5.3 1.5–12	1.8 ± 0.3 1.4–2.1
CB 114	1.5 ± 0.8 0.8–2.4	1.6 ± 0.7 0.5–3.6	4.2 n.d.–4.2	n.d.	0.2 ± 0.1 n.d.–0.2	0.2 n.d.–0.2	0.7 ± 0.2 n.d.–0.9	0.3 n.d.–0.3
CB 118	80 ± 47 35–128	47 ± 13 19–97	23 ± 16 14–41	6.5 ± 1.2 5.5–7.8	17 ± 3.1 13–19	6.0 ± 1.6 4.2–7.2	25 ± 19 7.5–46	5.5 ± 1.9 3.3–6.8
CB 123	14 ± 8.7 4.4–22	5.9 ± 2.2 2.7–12	3.3 ± 2.3 1.4–5.8	0.8 ± 0.3 n.d.–1.0	2.3 ± 0.5 1.8–2.7	0.7 ± 0.3 0.3–0.9	4.3 ± 3.9 0.8–8.5	0.7 0.4–1.0
CB 156	8.5 ± 4.7 3.3–12	2.3 ± 0.2 2.1–2.5	2.8 ± 2.1 1.0–5.4	0.8 ± 0.5 0.5–1.4	2.3 ± 0.8 1.5–3.0	0.6 ± 0.1 n.d.–0.6	3.3 ± 2.4 1.9–6.1	0.7 ± 0.3 n.d.–1.0
CB 157	1.7 ± 1.3 0.5–3.0	0.4 ± 0.1 0.3–0.4	n.d.	0.1 n.d.–0.1	1.8 ± 0.4 0.3–2.9	0.1 n.d.–0.1	1.0 ± 0.6 n.d.–1.4	0.6 n.d.–1.1
CB 167	5.1 ± 3.7 1.3–8.6	0.7 ± 0.2 0.6–0.9	2.8 ± 1.7 1.2–5.9	0.3 ± 0.2 n.d.–0.5	1.2 ± 0.4 0.5–2.0	0.3 ± 0.1 n.d.–0.4	1.6 ± 1.2 0.4–2.8	0.6 n.d.–0.6
CB 189	2.5 ± 0.3 n.d.–2.8	1.0 ± 0.7 n.d.–1.5	3.1 n.d.–3.1	n.d.	0.5 n.d.–0.5	0.1 n.d.–0.1	1.4 n.d.–1.4	n.d.
Σ mono-ortho PCBs	136 58–213	66 34–123	41 20–67	10 7.8–12	29 21–34	9.7 6.1–12	44 12–80	10 5.9–12
Σ_{12} dl-PCBs	143 66–219	71 37–126	41 20–67	11 9.0–13	32 24–37	11 7.1–11	47 12–83	12 7.0–13
Σ_{18} PCBs	779 429–1123	229 201–275	439 308–641	106 66–137	274 168–343	93 72–135	251 124–404	80 39–111
WHO ₂₀₀₅ -TEQs	0.16 0.07–0.22	0.03 0.02–0.03	0.15 0.01–0.39	0.06 0.01–0.15	0.10 0.03–0.20	0.04 0.02–0.07	0.07 0.04–0.09	0.06 0.02–0.11

The number of fish muscle homogenates per species per station (N) = 3; n.d. – non-detectable.

(such as CB 138 and CB 153) have lower chemical degradation rates than the less chlorinated PCBs and thus are retained in aquatic systems to a greater degree, where they may bioaccumulate in organisms like fish (Lavandier et al., 2013).

The PCB levels at station 1 were significantly higher ($p < 0.05$, Mann–Whitney U -test) than those from stations 3 and 4. However, statistical analysis between stations 1 and 2 showed no significant difference between the two locations. Station 1 is located near the mouth of the Nakivubo channel which brings in effluents from a number of chemical manufacturing factories, industrial waste treatment plants and municipal solid waste incinerators. Furthermore, in close vicinity of the station 1 are industrial facilities and Port bell where ships anchor on their way to and from Ports like Mwanza in Tanzania, Kisumu in Kenya and Jinja Port in Uganda. Although, the shipping industry promotes economic development in the area, it inevitably brings pollution related problems.

High concentrations of PCBs were also observed at station 2 (mean value of the Σ_{18} PCBs in sediments was 2576, range 1753–3970 pg g^{-1} dw). Station 2 is situated close to the Ggaba landing site and a local market. Besides the market a lot of commercial, domestic and industrial activities occur near the station and could be releasing domestically-produced commercial PCB mixtures into the lake. The PCB levels in sediments from the outer part of the Murchison Bay (stations 3 and 4), both far from industrial complexes, were an order of magnitude lower compared to those in the inner Bay (Fig. 3). The

low PCB levels in the outer Murchison Bay are likely due to the far offshore distance from the pollution sources.

The WHO₂₀₀₅-TEQs for the 12 dl-PCBs in sediments of the present study varied from 0.04 to 0.64 pg TEQ g^{-1} . Station 1 had the highest TEQ values ranging from 0.37 to 0.64 pg TEQ g^{-1} (mean 0.55 pg TEQ g^{-1}) while station 4 had the lowest (range 0.07 to 0.14, mean 0.10 pg TEQ g^{-1}). CB 126 was the highest contributor to the Σ_{12} TEQs (70–89%) compared to other congeners at all locations. The WHO₂₀₀₅-TEQs in most samples from the Murchison Bay were below the interim sediment quality guideline of 0.85 pg TEQ g^{-1} dw recommended by the Canadian Council of Ministers of the Environment (2002). Thus the dl-PCB concentrations in the present work are unlikely to pose a threat to benthic organisms.

3.2. Concentrations of PCBs in fish from the Murchison Bay

Table 3 presents the PCB concentrations in fish muscle homogenates of *L. niloticus* and *O. niloticus*. Average concentrations of the Σ_{18} PCBs ranged from 80 to 779 pg g^{-1} wet weight (ww). The concentrations were in the same range as those reported in similar fish species from the Napoleon Gulf of Lake Victoria (94–716 pg g^{-1} ww: Ssebugere et al., 2013a; Ssebugere et al., 2014) and River Nile in Egypt (695–853 pg g^{-1} ww; El-Kady et al., 2007). In neighboring Burundi, Manirakiza et al. (2002) reported PCB concentrations between 44900 and 166700 pg g^{-1} lw in *O. niloticus* collected from Lake Tanganyika.

Fig. 4. Mean PCB concentrations in fish per station. a) Mean PCB concentrations in *L. niloticus* per station. b) Mean PCB concentrations in *O. niloticus* per station.

These levels were at least an order of magnitude higher compared to the present results. In Egypt, Said et al. (2008) reported PCB concentrations varying from 3320 to 72060 pg g^{-1} lw in fish (*O. niloticus* and *Clarias gariepinus*) from Lake Burullus. The concentrations were again higher than the data in the study herein. Studies outside Africa have also reported PCBs in fish. In Hawai'i, PCB levels ranging from 51900 to 89420 pg g^{-1} lw were reported in tilapia (*Oreochromis mossambicus*) collected from the Manoa stream and Ala Wai Canal of O'ahu (Yang et al., 2008). The PCB levels were also higher than those reported in this study. In South China, Nie et al. (2006) reported PCB concentrations ranging from 5150 to 226000 pg g^{-1} lw in tilapia (*Tilapia mossambica*) from Pearl River Delta. The concentrations are still higher compared to the data in the current study.

The $\Sigma_{18}\text{PCBs}$ in *L. niloticus* at station 1 was one-fold higher than that at station 2 and about three times higher than that at station 4. The $\Sigma_{18}\text{PCBs}$ at stations 3 and 4 were quite similar. For *O. niloticus*, still station 1 showed the highest levels of the $\Sigma_{18}\text{PCBs}$ (mean value 229 pg g^{-1}) compared to other stations. Station 1 was followed by station 2 and then station 3, while station 4 had the lowest concentration. The concentrations of the different PCB congeners at the different stations for the 2 fish species are shown in Fig. 4.

The PCB concentrations in *L. niloticus* were markedly higher ($p < 0.05$, Mann–Whitney U -test) than those in *O. niloticus* at all study sites. The nature of the diet could be a possible explanation for such an inter-specific differences in concentrations. *L. niloticus* are exclusively piscivorous, consuming all available fish species including its

own siblings whereas *O. niloticus* relies on zooplankton and macro-invertebrates (Ogari, 1984; Njiru et al., 2004). Earlier studies established that PCBs can have great variations among different fish, as well as within species depending on their feeding habits and trophic levels (Suedel et al., 1994; Vander Zanden and Rasmussen, 1996; Davis et al., 2002; van der Oost et al., 2003).

On the whole, the Σ_6 indicator PCBs in fish of the present study were an order of magnitude higher than the Σ_7 mono-ortho-PCBs and Σ_4 non-ortho-PCBs. However, the Σ_6 indicator PCB levels (83–636 pg g^{-1}) in this study were within the limit of 75000 pg g^{-1} ww set for fish by the European Commission (EC, 2011), implying that the study fish were fit for human consumption in regard to indicator PCBs. Indicator PCB congeners such as CB 138 and CB 153 which were predominant in the sediments were also major contributors in *L. niloticus* (Fig. 5). However, variations in contribution of the indicator PCBs to the $\Sigma_{18}\text{PCBs}$ were observed for *L. niloticus* at the different stations. For the coplanar PCBs, the mono-ortho PCBs were the predominant congeners in most fish (their contribution to the $\Sigma_{12}\text{PCBs}$ ranged from 84 to 100%). CB 118 was the most abundant congener, accounting for 46–69% of the $\Sigma_{12}\text{PCBs}$, in comparison to other coplanar compounds. The high levels of CB 118 in the two fish species could be due to the inability of the species to metabolize the congener after accumulation. The presence of CB 118 in our samples suggests paint additives, municipal waste plants, iron ore sintering plants or medical waste incinerators as potential sources of PCBs (Abad et al., 2006; Aries et al., 2006; Shin et al., 2006; Chen et al., 2009; Jartun et al., 2009). Furthermore, it is likely that the

Fig. 5. Percent contribution of the congeners to the ΣPCBs in the fish. a) Contribution of the different congeners to the $\Sigma_{18}\text{PCBs}$ in *L. niloticus*. b) Contribution of the different congeners to the $\Sigma_{18}\text{PCBs}$ in *O. niloticus*.

PCBs came from the residues of former applications like transformers, electrical equipment and other industrial uses. The use of these applications containing PCBs was common in Uganda before their commercial production was banned worldwide in 1979 (National Environment Management Authority, 2007).

The WHO₂₀₀₅-TEQs for dl-PCBs in the study herein were 0.01–0.39 and 0.01–0.11 pg g⁻¹ ww for *L. niloticus* and *O. niloticus*, respectively. The non-ortho PCBs exhibited the highest contribution to the Σ₁₂PCBs (>75%) compared to the mono-ortho PCBs in both fish species. A similar observation was also reported by Ssebugere et al. (2013a) in fish from the Napoleon Gulf and Thurston Bay of Lake Victoria. The WHO₂₀₀₅-TEQs in the present study were within the permissible level of 3.5 pg g⁻¹ ww set by the European Commission (EC, 2011), suggesting that the fish from the Murchison Bay were safe for consumption with respect to dl-PCBs.

4. Conclusions

To our knowledge this is the first study to record PCB concentrations in sediments and fish from the Murchison Bay. The station closest to Nakivubo channel had the highest PCB concentrations compared to other locations. Land based activities such as urban centers, commercial and industrial establishments are likely the major sources of PCB contamination. The PCB concentrations in the present study were generally comparable to those of most studies in Africa but lower than the data examined elsewhere in the world. The WHO-TEQs were below the recommended values set for sediments and fish, indicating that the risk posed by PCBs is limited, compared to other health and environmental issues, such as the high disease burden due to diarrhea and other infectious diseases, associated with a limited availability of sanitation and treated water (Yeka et al., 2012; Tumwebaze et al., 2013; McElligott et al., 2013; Bwire et al. 2013; Katukiza et al., 2014). However, regulatory controls for monitoring and mitigating wastewater emissions into Lake Victoria need to be implemented and emphasized by the relevant governmental agencies in Uganda.

Conflict of interest

There is no conflict of interest.

Acknowledgments

Patrick Ssebugere is grateful for the research fellowship by the Third World Academy of Sciences (TWAS) and Chinese Academy of Sciences (CAS). The authors would like acknowledge the financial support from the Chinese Academy of Sciences (YSW2013B01) and National Natural Science Foundation of China (41276195, 21277165 and 21107121).

References

- Abad E, Martinez K, Caixach J, Rivera J. Polychlorinated dibenzo-p-dioxins, dibenzofurans and 'dioxin-like' PCBs in flue gas emissions from municipal waste management plants. *Chemosphere* 2006;63:570–80.
- Aberg A, Macleod M, Wiberg K. Physical–chemical property data for dibenzo-p-dioxins (DD), dibenzofuran (DF), and chlorinated DD/Fs: a critical review and recommended values. *J Phys Chem Ref Data* 2008;37:1997–2008.
- Aberg A, Tysklind M, Nilsson T, Macleod M, Hanberg A, Andersson R, et al. Exposure assessment at a PCDD/F contaminated site in Sweden—field measurements of exposure media and blood serum analysis. *Environ Sci Pollut Res* 2010;17:26–39.
- Adu-Kumi S, Kawano M, Shiki Y, Yeboah PO, Carboo D, Pwamang J, et al. Organochlorine pesticides (OCPs), dioxin-like polychlorinated biphenyls (dl-PCBs), polychlorinated dibenzo-p-dioxins and polychlorinated dibenzo furans (PCDD/Fs) in edible fish from Lake Volta, Lake Bosomtwi and Weija Lake in Ghana. *Chemosphere* 2010;81(6):675–84.
- Ameur WB, Trabelsi S, El Bedoui B, Driss MR. Polychlorinated biphenyls in sediments from Ghar El Melh lagoon, Tunisia. *Bull Environ Contam Toxicol* 2011;86(5):539–44.
- Aries E, Anderson DR, Fisher R, Fray TAT, Hemfrey D. PCDD/F and "dioxin-like" PCB emissions from iron ore sintering plants in the UK. *Chemosphere* 2006;65:1470–80.

- Banadda EN, Kansime F, Kigobe M, Kizza M, Nhapi I. Land use based non-point source pollution: a threat to Murchison bay water quality in Uganda. *Water Qual II Suppl* 2009;1:93–104.
- Banadda N, Ayaa F, Nhapi I, Wali UG, Kimwaga RJ, Mashauri DA. Modeling diffusive flux of non point source pollutants in Lake Victoria: a comparison study of Fick's law and the Fokker–Planck law. *Open Environ Eng J* 2011a;4:105–11.
- Banadda N, Nhapi I, Wali UG. Determining and modeling the dispersion of non point source pollutants in Lake Victoria: a case study of Ggaba landing site in Uganda. *Afr J Environ Sci Technol* 2011b;5(3):178–85.
- Barakat AO, Mostafa A, Wade TL, Sweet ST, El Sayed NB. Spatial distribution and temporal trends of persistent organochlorine pollutants in sediments from Lake Maryut, Alexandria, Egypt. *Mar Pollut Bull* 2012;64(2):395–404.
- Bwire G, Malimbo M, Maskery B, Kim EY, Mogasale V, Levin A. The burden of cholera in Uganda. *PLoS Negl Trop Dis* 2013;7(12):e2545.
- Canadian Council of Ministers of the Environment. Canadian environmental quality guidelines for polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans (PCDD/Fs). Technical supporting document. Ottawa: National Guidelines and Standards Office, Environmental Quality Branch; 2002.
- Carvalho FP, Villeneuve JP, Cattini C, Tolosa I, Thuan DD, Nhan DD. Agrochemical and polychlorobiphenyl (PCB) residues in the Mekong River delta, Vietnam. *Mar Pollut Bull* 2008;56(8):1476–85.
- Castells P, Parera J, Santos FJ, Galceran MT. Occurrence of polychlorinated naphthalenes, polychlorinated biphenyls and short-chain chlorinated paraffins in marine sediments from Barcelona (Spain). *Chemosphere* 2008;70(9):1552–62.
- Chen T, Li XD, Yan JH, Jin YQ. Polychlorinated biphenyls emission from a medical waste incinerator in China. *J Hazard Mater* 2009;172:1339–43.
- Covaci A, Gheorghe A, Voorspoels A, Maervoet J, Redeker ES, Blust R, et al. Polybrominated diphenyl ethers, polychlorinated biphenyls and organochlorine pesticides in sediment cores from Western Scheldt River (Belgium): analytical aspects and depth profiles. *Environ Int* 2005;31:367–75.
- Darko G, Akoto O, Oppong C. Persistent organochlorine pesticide residues in fish, sediments and water from Lake Bosomtwi, Ghana. *Chemosphere* 2008;72(1):21–4.
- Davis JA, May MD, Greenfield BK, Fairey R, Roberts C, Ichikawa G, et al. Contaminant concentrations in sport fish from San Francisco Bay, 1997. *Mar Pollut Bull* 2002;44:1117–29.
- EC. Commission Regulation (EU) No 1259/2011 of 2 December 2011 amending regulation (EC) No 1881/2006 as regards maximum levels for dioxins, dioxin-like PCBs and non dioxin-like PCBs in foodstuffs. *Off J Eur Union* 2011;320/18–23.
- El-Kady AA, Abdel-Wahhab MA, Henkelmann B, Belal MH, Morsi MK, Galal SM, et al. Polychlorinated biphenyl, polychlorinated dibenzo-p-dioxin and polychlorinated dibenzofuran residues in sediments and fish of the River Nile in the Cairo region. *Chemosphere* 2007;68(9):1660–8.
- Eqani SA, Malik RN, Cincinelli A, Zhang G, Mohammad A, Qadir A, et al. Uptake of organochlorine pesticides (OCPs) and polychlorinated biphenyls (PCBs) by river water fish: the case of River Chenab. *Sci Total Environ* 2013;450–451:83–91.
- Gidararakos E, Petrantonaki M, Anastasiadou K, Schramm KW. Characterization and hazard evaluation of bottom ash produced from incinerated hospital. *J Hazard Mater Waste* 2009;172(2–3):935–42.
- Gustavson KE, Burton GA, Francingues NR, Reible DD, Vorhees DJ, Wolfe JR. Evaluating the effectiveness of contaminated-sediment dredging. *Environ Sci Technol* 2008;42:5042–7.
- Ha MH, Lee DH, Jacobs DR. Association between serum concentrations of persistent organic pollutants and self-reported cardiovascular disease prevalence. Results from the national health and nutrition examination survey, 1999–2002. *Environ Health Perspect* 2007;115:1204–9.
- Ha MH, Lee DH, Son HK, Park SK, Jacobs DR. Association between serum concentrations of persistent organic pollutants and prevalence of newly diagnosed hypertension: results from the National Health and Nutrition Examination Survey 1999–2002. *J Hum Hypertens* 2009;23:274–86.
- Haande S, Rohrlack T, Semyalo PR, Brettum P, Edvardsen B, Lyche-Solheim A, et al. Phytoplankton dynamics and cyanobacterial dominance in Murchison Bay of Lake Victoria (Uganda) in relation to environmental conditions. *Limnologia* 2011;41:20–9.
- Hellar-Kihampa H, De Wael K, Lugwisha E, Malarvannan G, Covaci A, Van Grieken R. Spatial monitoring of organohalogen compounds in surface water and sediments of a rural–urban river basin in Tanzania. *Sci Total Environ* 2013;447:186–97.
- Helm PA, Gewurtz SB, Whittle DM, Marvin CH, Fisk AT, Tomy GT. Occurrence and biomagnification of polychlorinated naphthalenes and non- and mono-ortho PCBs in Lake Ontario sediment and biota. *Environ Sci Technol* 2008;42(4):1024–31.
- Huang X, Hites RA, Foran JA, Hamilton C, Knuth BA, Schwager SJ, et al. Consumption advisories for salmon based on risk of cancer and non cancer health effects. *Environ Res* 2006;101:263–74.
- Ilyas M, Sudaryanto A, Setiawan IE, Riyadi AS, Isobe T, Takahashi S, et al. Characterization of polychlorinated biphenyls and brominated flame retardants in sediments from riverine and coastal waters of Surabaya, Indonesia. *Mar Pollut Bull* 2011;62(1):89–98.
- Jartun M, Ottesen RT, Steinnes E, Volden T. Painted surfaces—important sources of polychlorinated biphenyls (PCBs) contamination to the urban and marine environment. *Environ Pollut* 2009;157(1):295–302.
- Jiao L, Zheng GJ, Minh TB, Richardson B, Chen L, Zhang Y, et al. Persistent toxic substances in remote lake and coastal sediments from Svalbard, Norwegian Arctic: levels, sources and fluxes. *Environ Pollut* 2009;157(4):1342–51.
- Jönsson A, Gustafsson Ö, Axelmann J, Sundberg H. Global accounting of PCBs in the continental shelf sediments. *Environ Sci Technol* 2003;37:245–55.
- Katukiza AY, Ronteltap M, van der Steen P, Foppen JWA, Lens PNL. Quantification of microbial risks to human health caused by waterborne viruses and bacteria in an urban slum. *J Appl Microbiol* 2014;116(2):447–63.

- Lavandier R, Quinete N, Hauser-Davis RA, Dias PS, Taniguchi S, Montone R, et al. Polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) in three fish species from an estuary in the southeastern coast of Brazil. *Chemosphere* 2013;90(9):2435–43.
- Lee DH, Lee IK, Jin SH, Steffes M, Jacobs Jr DR. Association between serum concentrations of persistent organic pollutants and insulin resistance among nondiabetic adults: results from the national health and nutrition examination survey 1999–2002. *Diabetes Care* 2007a;30:622–8.
- Lee DH, Lee IK, Porta M, Steffes M, Jacobs Jr DR. Relationship between serum concentrations of persistent organic pollutants and the prevalence of metabolic syndrome among non-diabetic adults: results from the National Health and Nutrition Examination Survey 1999–2002. *Diabetologia* 2007b;50:1841–51.
- Lundin L, Gomez-Rico MF, Forsberg C, Nordenskjöld C, Jansson S. Reduction of PCDD, PCDF and PCB during co-combustion of biomass with waste products from pulp and paper industry. *Chemosphere* 2013;91:797–801.
- Manirakiza P, Covaci A, Nizigiyimana L, Ntakimazi G, Schepens P. Persistent chlorinated pesticides and polychlorinated biphenyls in selected fish species from Lake Tanganyika, Burundi, Africa. *Environ Pollut* 2002;117:447–55.
- Martinez A, Norström K, Wang K, Hornbuckle CK. Polychlorinated biphenyls in the surficial sediment of Indiana Harbor and Ship Canal, Lake Michigan. *Environ Int* 2010;36(8):849–54.
- Mazet A, Keck G, Berny P. Concentrations of PCBs, organochlorine pesticides and heavy metals (lead, cadmium, and copper) in fish from the Drome river: potential effects on otters (*Lutra lutra*). *Chemosphere* 2005;61:810–6.
- McElligott JT, Naaktgeboren C, Makuma-Massa H, Summer AP, Deal JL. Prevalence of intestinal protozoa in communities along the Lake Victoria region of Uganda. *Int J Infect Dis* 2013;17(8):e658–9.
- Miller A, Hedman JE, Nyberg E, Haglund P, Cousins IT, Wiberg K, et al. Temporal trends in dioxins (polychlorinated dibenzo-*p*-dioxin and dibenzofurans) and dioxin-like polychlorinated biphenyls in Baltic herring (*Clupea harengus*). *Mar Pollut Bull* 2013;73:220–30.
- Mohammed A, Peterman P, Echols K, Feltz K, Tegerdine G, Manoo A, et al. Polychlorinated biphenyls (PCBs) and organochlorine pesticides (OCPs) in harbor sediments from Sea Lots, Port-of-Spain, Trinidad and Tobago. *Mar Pollut Bull* 2011;62(6):1324–32.
- Moon HB, Choi M, Choi HG, Kannan K. Severe pollution of PCDD/Fs and dioxin-like PCBs in sediments from Lake Shihwa, Korea: tracking the source. *Mar Pollut Bull* 2012;64:2357–63.
- National Environment Management Authority. Identification and quantification of dioxins and furans in Uganda: POPs inventory report. Uganda: Kampala; 2007.
- Nie PX, Lan YC, An CT, Wong HM, Li BK. Distributions and congener patterns of PCBs in fish from major aquaculture areas in the Pearl River Delta, South China. *Hum Ecol Risk Assess* 2006;12(2):363–73.
- Njiru M, Okeyo-Owuor BJ, Muchiri M, Cowx GI. Shifts in the food of Nile tilapia, *Oreochromis niloticus* (L.) in Lake Victoria, Kenya. *Afr J Ecol* 2004;42:163–70.
- Ogari J. Distribution, food and feeding habits of “*Lates niloticus*” in Nyanza Gulf of Lake Victoria (Kenya). *FAO Fish Rep* 1984;335:68–80.
- Parnell PE, Groce AK, Stebbins TD, Dayton PK. Discriminating sources of PCB contamination in fish on the coastal shelf off San Diego, California (USA). *Mar Pollut Bull* 2008;56:1992–2002.
- Said TO, El Moselhy KM, Rashad AA, Shreadah MA. Organochlorine contaminants in water, sediment and fish of Lake Burullus, Egyptian Mediterranean Sea. *Bull Environ Contam Toxicol* 2008;81(2):136–46.
- Shin SK, Kim KS, You JC, Song BJ, Kim JG. Concentration and congener patterns of polychlorinated biphenyls in industrial and municipal waste incinerator flue gas, Korea. *J Hazard Mater* 2006;133:53–9.
- Ssebugere P, Kiremire BT, Henkelmann B, Bernhöft S, Wasswa J, Kasozi GN, et al. PCDD/Fs and dioxin-like PCBs in surface sediments from Lake Victoria, East Africa. *Sci Total Environ* 2013a;454–455:528–33.
- Ssebugere P, Kiremire TB, Henkelmann B, Bernhöft S, Kasozi GN, Wasswa J, Schramm K-W. PCDD/Fs and dioxin-like PCBs in fish species from Lake Victoria, East Africa. *Chemosphere* 2013b;92(3):317–21.
- Ssebugere P, Sillanpää M, Kiremire TB, Gabriel Kasozi NK, Wang P, Samuel Sojinu OS, Otieno OP, Zhu N, Zhu C, Zhang H, Shang H, Ren D, Li Y, Zhang Q, Jiang G. Polychlorinated biphenyls and hexachlorocyclohexanes in sediments and fish species from the Napoleon Gulf of Lake Victoria, Uganda. *Sci Total Environ* 2014;481:55–60.
- Suedel BC, Boraczek JA, Peddicord RK, Clifford PA, Dillon TM. Trophic transfer and biomagnification potential of contaminants in aquatic ecosystems. *Rev Environ Contam Toxicol* 1994;136:21–89.
- Sundqvist KL, Tysklind M, Cato I, Bignert A, Wiberg K. Levels and homologue profiles of PCDD/Fs in sediments along the Swedish coast of the Baltic Sea. *Environ Sci Pollut Res* 2009a;16:396–409.
- Sundqvist KL, Tysklind M, Geladi P, Cato I, Wiberg K. Congener fingerprints of tetra-through octa-chlorinated dibenzo-*p*-dioxins and dibenzofurans in Baltic surface sediments and relations to potential sources. *Chemosphere* 2009b;77:612–20.
- Tumwebaze IK, Orach CG, Niwagaba C, Luthi C, Mosler HJ. Sanitation facilities in Kampala slums, Uganda: users' satisfaction and determinant factors. *Int J Environ Health Res* 2013;23(3):191–204.
- Uemura H, Arisawa K, Hiyoshi M, Kitayama A, Takami H, Sawachika F, et al. Prevalence of metabolic syndrome associated with body burden levels of dioxin and related compounds among Japan's general population. *Environ Health Perspect* 2009;117:568–73.
- Van den Berg M, Birnbaum LS, Denison M, De Vito M, Farland W, Feeley M, et al. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci* 2006;93:223–41.
- Van der Oost R, Beyer J, Vermeulen NPE. Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environ Toxicol Pharmacol* 2003;13:57–149.
- Vander Zanden MJ, Rasmussen JB. A trophic position model of pelagic food webs: impact on contaminant bioaccumulation in lake trout. *Ecol Monogr* 1996;66:451–77.
- Vane HC, Harrison I, Kim WA. Polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) in sediments from the Mersey Estuary, U.K. *Sci Tot Environ* 2007;374:112–26.
- Verhaert V, Covaci A, Bouillon S, Abrantes K, Musibono D, Bervoets L, et al. Baseline levels and trophic transfer of persistent organic pollutants in sediments and biota from the Congo River Basin (DR Congo). *Environ Int* 2013;59:290–302.
- Vives I, Canuti E, Castro-Jiménez J, Christoph EH, Eisenreich SJ, Hanke G, et al. Occurrence of polychlorinated dibenzo-*p*-dioxins and dibenzofurans (PCDD/Fs), polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) in Lake Maggiore (Italy and Switzerland). *J Environ Monit* 2007;9(6):589–98.
- Wang P, Zhang H, Fu J, Li Y, Wang T, Wang Y, et al. Temporal trends of PCBs, PCDD/Fs and PBDEs in soils from an E-waste dismantling area in East China. *Environ Sci Process Impacts* 2013;15(10):1897–903.
- Wasswa J. Characterization of selected pesticide residues in sediments of the Ugandan side of Lake Victoria. PhD Thesis Kampala, Uganda: Department of Chemistry, Makerere University; 2009.
- Wasswa J, Kiremire BT, Nkedi-Kizza P, Mbabazi J, Ssebugere P. Organochlorine pesticide residues in sediments from the Ugandan side of Lake Victoria. *Chemosphere* 2010;82(1):130–6.
- Wiberg K, McLachlan M, Jonsson P, Johansson N, Josefsson S, Knekt E, et al. Sources, transport reservoirs and fate of dioxins, PCBs, and HCB in the Baltic Sea environment. Report No. 5912 for the Swedish Environmental Protection Agency 978-91-620-5912-5; 2009.
- Yang F, Wilcox B, Jin S, Alonso Aguirre A, Rougée L, Xu Y, Lu Y. Detection and quantitative analysis of polychlorinated biphenyls in tilapia from Hawaiian waters. *Chemosphere* 2008;73(1):133–7.
- Yeka A, Gasasira A, Mpimbaza A, Achan J, Nankabirwa J, Nsobya S, et al. Malaria in Uganda: challenges to control on the long road to elimination: I. Epidemiology and current control efforts. *Acta Trop* 2012;121(3):184–95.